

**THE ALLEVIATION OF SALINITY INDUCED STRESS WITH THE APPLICATION
OF SILICON IN SOILLESS GROWN *Lactuca sativa* L. 'Eish!'**

by

CHRISTOPHER JODI MILNE

Thesis submitted in fulfilment of the requirements for the degree

Master of Technology: Horticulture

in the Faculty of Applied Sciences

at the Cape Peninsula University of Technology

Supervisor: Prof. CP Laubscher

Co-supervisor: Prof. PA Ndakidemi

**Cape Town
June 2012**

DECLARATION

I, Christopher Jodi Milne, declare that the contents of this dissertation/thesis represent my own unaided work, and that the dissertation/thesis has not previously been submitted for academic examination towards any qualification. Furthermore, it represents my own opinions and not necessarily those of the Cape Peninsula University of Technology.

Signed

Date

ABSTRACT

This article based thesis includes two individual studies evaluating the role of silicon (Si) in mitigating the negative effects that are associated with sodium chloride (NaCl) induced toxicity in lettuce (*Lactuca sativa* L. 'Eish!').

The objectives of the studies were to evaluate the effects Si on NaCl stressed lettuce. This was done by assessing yield parameters, including fresh and dry root and shoot weights, chlorophyll content, Si, sodium (Na) and chlorine (Cl) concentrations in plant tissues, total polyphenolics (TP), oxidative radical absorbance capacity (ORAC), thiobarbituric acid reactive substance (TBARS), catalase (CAT), glutathione (GSH), and oxidised glutathione (GSSG). NaCl was applied at two different concentrations; 30 and 60 mM, and Si at four different concentrations; 0, 1, 2 and 4 mM.

Notable results include a significant increase in fresh root and shoot weight with additions of 2 mM Si compared to that of the 60 mM NaCl control, with notable, yet insignificant increases with Si concentrations of 1 and 4 mM. Dry root and shoot weight with additions of 1 mM Si, and in shoot weight with additions of 2 mM Si, significantly increased when compared to the 60 mM NaCl control. Increases in dry root and shoot weight of the other Si additions lead to clear, yet insignificant increases in dry weights. Few differences in Cl content were evident amongst treatments, besides a significant increase in root content and a significant decrease in shoot content with applications of 4 mM Si exposed to 60 mM NaCl. Na shoot content of 1, 2 and 4 mM Si at 60 mM NaCl showed a significant decrease compared to that of the control. Applications of 2 and 4 mM Si showed increased shoot Si when compared to both the 30 and 60 mM NaCl controls.

Other findings included increases in ORAC values, significantly with 4 mM Si at both 30 and 60 mM NaCl. Additionally, GSH concentrations significantly increased in 2 mM Si treated plants when compared to the 30 mM NaCl control, with increases in both GSH and GSSG at 1 and 4 mM Si treated plants when compared to the 60 mM NaCl control. CAT activity decreased with applications of Si, significantly with 1 and 4 mM NaCl when compared to the 30 mM control. TP concentrations stayed constant throughout treatments, with only marginal differences between the two controls.

It can be concluded that the applications of Si is of definite benefit to lettuce plant growth. These benefits are especially evident at higher NaCl concentrations. Although the regular inclusion of Si in nutrient solutions can be recommended at the NaCl concentrations that

were used in this study, studies evaluating the effects of Si on lettuce growth at higher NaCl concentrations could be of value.

ACKNOWLEDGEMENTS

I wish to thank:

Prof. Charles Laubscher, my supervisor and Prof. Patrick Ndakidemi, my co-supervisor, for advice during the experimental and writing process.

Mrs Corrie Uys for her help with the statistics and always being available to assist even at the shortest notice.

Prof. Jeanine Marnewick for her time, guidance and support with regards to antioxidants and oxidative stress, especially during the writing of this study.

Olatogni Berenice Alinde and Fanie Rautenbach for their professionalism, friendliness, and exceptional patience while training me in laboratory procedures and for sharing their expertise, especially with regards to assays and mathematical calculations.

The Mauerberger Foundation for granting me the Mauerberger Foundation Scholarship, allowing me to focus solely on this thesis.

The Cape Peninsula University of Technology for the financial assistance, allowing me to purchase the resources that made these studies possible.

Finally, I wish to thank my father, mother and sister, Glenn, Sharon and Chantal Milne, my great friend Geert Sprangers and my partner Güler Aslan, for their continual help, support and motivation, without which this thesis would have never been successful.

Declaration	ii
Abstract	iii
Acknowledgements	v
Table of Contents	vi
List of Tables and Figures	viii
Appendices	x
Glossary	xi

CHAPTER ONE: INTRODUCTION	1
----------------------------------	----------

CHAPTER TWO: LITERATURE REVIEW

2.1	Salinity and its effects on plants	2 – 4
2.2	Silicon and salinity stress mitigation	4
2.3	Antioxidants and salinity stress	5 – 6
2.4	Antioxidants interactions with silicon and salinity stress	7 – 8

CHAPTER THREE: THE ALLEVIATION OF SALINITY INDUCED STRESS WITH APPLICATIONS OF SILICON IN SOILLESS GROWN *Lactuca sativa* L. ‘Eish!’

3.1	Abstract	12
3.2	Introduction	12 – 13
3.3	Materials and Methods	13 – 14
3.3.1	Basal Nutrient Solution	14
3.3.2	Silicon and NaCl Treatments	14
3.3.3	Plant Growth Parameters	15
3.3.4	Determination of Chlorophyll Content in Plant Leaves	15
3.3.5	Elemental Analysis	15 – 16
3.3.6	Statistical Analysis	16
3.4	Results	16 – 18
3.5	Discussion	18 – 19
3.6	Acknowledgments	19

CHAPTER FOUR: SALINITY INDUCED CHANGES IN OXIDATIVE STRESS AND ANTIOXIDANT STATUS AS AFFECTED BY APPLICATIONS OF SILICON IN LETTUCE (*Lactuca sativa* L. 'Eish!').

4.1	Abstract	32
4.2	Introduction	32 – 33
4.3	Materials and Methods	34
4.3.1	Plant Conditions and Cultivation	34
4.3.2	Basal Nutrient Solution	34
4.3.3	Silicon and NaCl Treatments	34 – 35
4.3.4	Sampling and Harvesting	35
4.3.4.1	Antioxidant Status and Oxidative Damage	35
4.3.4.2	Plant Yield and Elemental Analysis	35
4.3.5	Total Protein Analysis	35 – 36
4.3.6	Polyphenol Antioxidant Content and Capacity	36
4.3.6.1	Total Polyphenol Analysis	36
4.3.6.2	Oxygen Radical Absorbance Capacity Analysis	36
4.3.7	Catalase Analysis	36 – 37
4.3.8	Glutathione Analysis	37
4.3.9	Lipid Peroxidation Analysis	37 – 38
4.3.10	Statistical Analysis	38
4.4	Results	38 – 39
4.5	Discussion	40 – 41
4.6	Acknowledgments	42
4.7	References	42 – 47

CHAPTER FIVE: GENERAL DISCUSSION AND CONCLUSION 55 – 56

REFERENCES 57 – 64

LIST OF FIGURES

CHAPTER 2

- Figure 2.1:** Gallic Acid **9**
- Figure 2.2:** The ascorbate-glutathione cycle. **10**
- Figure 2.3:** Lipid peroxidation. **11**

CHAPTER 3

- Figure 3.1:** The effect of 30 and 60 mM NaCl on plant height, with additions of 0,1, 2 and 4 mM Si. Measurements were take weekly from 27 May (1) to 17 June (4) 2011 **23**
- Figure 3.2:** The effect of 30 and 60 mM NaCl on fresh root weight, with additions of 0, 1, 2 and 4 mM Si **24**
- Figure 3.3:** The effect of 30 and 60 mM NaCl on fresh shoot weight, with additions of 0, 1, 2 and 4 mM Si. **25**
- Figure 3.4:** The effect of 30 and 60 mM NaCl on dry root weight, with additions of 0, 1, 2 and 4 mM Si. **26**
- Figure 3.5:** The effect of 30 and 60 mM NaCl on dry shoot weight, with additions of 0, 1, 2 and 4 mM Si **27**
- Figure 3.6:** The effect of 30 and 60 mM NaCl on Na root content, with additions of 0, 1, 2 and 4 mM Si **28**
- Figure 3.7:** The effect of 30 and 60 mM NaCl on total chlorophyll content with additions of 0, 1, 2 and 4 mM Si. **29**

CHAPTER 4

Figure 4.1: Effect of Si applications on Si shoot concentrations in salinity stressed lettuce. **48**

Figure 4.2: Effect of Si applications on Oxidative Radical Absorbance Capacity (ORAC) values in salinity stressed lettuce. **49**

Figure 4.3: Effect of Si applications on Catalase (CAT) concentrations in salinity stressed lettuce. **50**

Figure 4.4: Effect of Si applications on Glutathione (GSH) concentrations in salinity stressed lettuce **51**

LIST OF TABLES

CHAPTER 3

Table 3.1: The effect of 30 and 60 mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4 mM Si. **30**

Table 3.2: The effect of 30 and 60 mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4 mM Si. **31**

CHAPTER 4

Table 4.1: Effects of Si applications on plant fresh and dry root and shoot weights in salinity stressed lettuce. **52**

Table 4.2: Effects of Si applications on Total Polyphenolics (TP) and Oxidative Radical Absorbance Capacity, in salinity stressed lettuce. **53**

Table 4.3: Effects of Si applications on Thiobarbituric Acid Reactive Substances (TBARS), Glutathione (GSH), Glutathione disulfide (GSSG), GSH:GSSG ratio and Catalase (CAT), in salinity stressed lettuce. **54**

APPENDICIES

APPENDIX A: The subirrigative, closed soilless growing system used for the cultivation of lettuce.

APPENDIX B: Paper published in The International Journal of the Physical Sciences.

APPENDIX C: Paper accepted by the International Journal of Agriculture and Biology.

GLOSSARY

Terms/Acronyms/Abbreviations	Definition/Explanation
Anion	Negatively charged ion
Antioxidant	A substance that hinders or prevents oxidation of an oxidizable substrate.
APX	Ascorbate peroxidase
Ascorbate-glutathione cycle	A biochemical pathway utilising glutathione, whereby ascorbate neutralises H_2O_2 and is recycled.
Cation	Positively charged ion
CAT	Catalase
Cl	Chlorine
EC	Electrical conductivity
GSSG	Glutathione disulphide (oxidised glutathione)
GSH	Glutathione
H_2O_2	Hydrogen peroxide
H_4SiO_4	Silicic acid
K	Potassium
Lipid peroxidation	Oxidative degradation of lipids.
mM	millimolar
$mS\ cm^{-1}$	milliSiemens per centimetre
MDA	Malondialdehyde
Na	Sodium
NaCl	Sodium chloride
O_2	Molecular oxygen
O_2^-	Superoxide
OH^{\cdot}	Hydroxyl ion
Opal	Amorphous silica
ORAC	Oxidative radical absorbance capacity
Oxidation	A loss of electrons by a chemical species.
Phenolics	Organic compounds comprised of a hydroxyl group bonded to an aromatic hydrocarbon.
Redox reaction	Oxidation-reduction reaction whereby electrons are transferred between species.

ROS	Reactive oxygen species
Salinisation	The accumulation of salts in a medium
Salt	Ionic compound that is a product of a neutralisation reaction.
Si	Silicon
SiO₂	Silica (Silicon dioxide)
TBARS	Thiobarbituric acid reactive substance
TP	Total polyphenolics

CHAPTER ONE

INTRODUCTION

This thesis is divided into five chapters: Chapter 1: Introduction, Chapter 2: Literature Review, Chapter 3: The alleviation of salinity induced stress with the applications of silicon in soilless grow *Lactuca sativa* L. 'Eish!', Chapter 4: Salinity induced changes in oxidative stress and antioxidant status as affected by applications of silicon in lettuce (*Lactuca sativa* L. 'Eish!'), Chapter 5: General Discussion and Conclusion, and Chapter 6: References. Chapter 3 and 4 are prepared in publishable format, and the layout is that of a journal article.

In addition to these foundation chapters, material that has been deemed relevant and complementary to this thesis, but incompatible with the articles, has been included at the end under the heading 'Appendices'.

Chapter 2 covers all aspects of salinity stress, silicon and antioxidants, the core subjects of this thesis and studies found herein. Salinity stress is introduced with the appreciation of the extent of salinity, the salts causing salinity, with the chemistry of this ancient problem being discussed in more detail in following paragraphs. The focal salt, NaCl, is then discussed, with the role of its cation Na⁺ and anion Cl⁻ in relation to plant nutrition and toxicity, examined.

The basic chemistry of Si is examined next, with the role of this element, especially with regards to its beneficial, ameliorative properties. An emphasis is placed on silicon's more recent quasi-essential status in plant nutrition, as opposed to its previous non-essential status. This discussion focuses on the ability of Si to mitigate salinity stress, specifically NaCl induced salinity stress, and the mechanisms that have been proposed by various authors.

ROS and antioxidants are discussed, centred on the antioxidants that were investigated in chapter 4, namely CAT, GSH (and GSSG) and polyphenolics. The chemistry, role and basic function of these antioxidants are discussed in a manner appropriate for a broad understanding. This understanding is further utilised to describe the relationship between stress, particularly salinity stress, antioxidants and ROS. Malondialdehyde (MDA), a product of ROS damage is discussed towards the close of the review. Chapter 2 concludes with the converging of the core subjects of this thesis; salinity stress, Si and antioxidants.

CHAPTER TWO

LITERATURE REVIEW

2.1 Salinity and its effects on plants

Salinity is defined as the quantity of NaCl, or as the quantity of total dissolved salts in a unit of water (Ghassemi et al., 1995). The dominant salts causing saline condition are chlorides, sulphates and bicarbonates of sodium, calcium and magnesium (Carter, 1981; Ghassemi et al., 1995). Of these salts, NaCl, as expressed by the definition, is the most common salt causing salinity (Harris, 1992:215; Epstein & Bloom, 2005:332-334).

Salts are found naturally in all soils (National Research Council, 1990), but the build-up of these salts in water and soil, known as salinisation (Ghassemi et al., 1995), is an immense problem - early stages of salinisation reduce soil productivity, and advanced stages destroy vegetation completely, thus ruining fertile land and diminishing biodiversity (Ghassemi et al., 1995).

Salinisation may be caused by long-term, natural processes, such as accumulation of weathered material, rain water containing dissolved solids or the appearance of the ocean over land (Ghassemi et al., 1995). These primary causes of salinisation are associated with conditions wherever evaporation exceeds precipitation (Epstein & Bloom, 2005:330-333). Salinisation, due to natural causes, is worldwide, occurring on all continents (Ghassemi *et al.*, 1995).

Saline conditions caused by human activities greatly contribute to the salinisation problem (Epstein & Bloom, 2005:330-333). This human, secondary salinisation, is caused by the mobilisation of salts stored in the soil or groundwater by the introduction of water through activities, like irrigation or land clearing (Ghassemi et al., 1995). The addition of water through practices, such as irrigation, increases the water table, and upon reaching the surface evaporates leaving the salts behind. The irrigation water itself will certainly contain dissolved solutes, substantially contributing to salinisation (Epstein & Bloom, 2005:330-333; Ghassemi et al., 1995). Saline conditions are by no means a recent problem, with the decline of the Mesopotamian civilizations associated with the salinisation of previously fertile land (Lerner et al., 1993:39). Since then, many factors have increased salinisation, including an increase in population leading to increased food demand, increases in water usage and a subsequent over utilisation of groundwater (Ghassemi et al., 1995).

With the increase in soluble salts dissolved in water there will be an increase in the ability of the water to conduct electricity (Resh, 1995:109). This electrical conductivity (EC) is measured using an EC device, with EC expressed in milliSiemens per cm (mS cm^{-1}). With an increase in the EC, there is an increase in the osmotic potential of the soil solution, and because osmosis is the principal way by which water enters plants (Stern, 2006:151-152) it will have a direct effect on plant-water availability (Carter, 1981). Resh (1995:109) states that salt levels above 4 mS cm^{-1} leads to wilting, suppressed growth and fruit cracking, with desired levels for plant growth between $2 - 4 \text{ mS cm}^{-1}$. This correlates with Ghassemi et al. (1995) who confirms that when a saturated soil extract exceeds 4 mS cm^{-1} , it is considered saline. Relatively high EC levels, leading to osmotic stress, is the primary cause of decreased growth and productivity in plants growing in saline conditions (Carter, 1981; Ghassemi et al., 1995). Although preferred EC levels are known, as stated by Resh (1995:109), tolerable EC can differ from one plant to the next (Maas & Hoffman, 1977; Shannon & Grieve, 1999). Above this salinity threshold level, there is a noted decrease in plant growth and size with an increase in salinity. Tolerable levels range, for example in strawberries, with a low threshold tolerance of 1 mS cm^{-1} , to cotton with a high threshold tolerance of 7.7 mS cm^{-1} (Maas & Hoffman, 1977). Observable symptoms due to osmotic stress usually include stunting of growth, the most common symptom, and dark green, thicker and more succulent leaves (Maas & Hoffman, 1977). Affected plants could also show the same symptoms as drought stress, due to the decreased availability of water (Carter, 1981).

As discussed earlier, NaCl is the most common salt causing saline conditions, leading to an abundance of available Na^+ and Cl^- ions. Cl is known to be an essential micronutrient in plants (Epstein & Bloom, 2005:210-211; Stern, 2006:161), and some argue that Cl should be classified as a macronutrient, contributing as much as 1% of the dry weight of some plants (Muckle, 1993). Na, on the other hand, is required by only a few plants (Epstein & Bloom, 2005:231-232). High concentrations of both of these ions are associated with leaf scorching (Shannon & Grieve, 1999), necrosis and defoliation (Maas & Hoffman, 1977). Additionally, excess NaCl has been associated with decreased photosynthesis and chlorophyll content (Moradi & Ismail, 2007; Savvas et al., 2009). Epstein & Bloom (2005:332-333), summarise the fundamental characteristics of salt-plant relations, and state that when Na is plentiful, above the micromolar range, it can displace and disrupt the functioning of K.

Plant species, cultivars and even individual plants of the same cultivar, differ in their tolerance to saline conditions (Rush & Epstein, 1978; Flowers & Yeo, 1981; Barroso & Alvarez, 1997). Lettuce, the crop used in this plant study, has an EC threshold tolerance of 1.3 mS cm^{-1} , and is subsequently classified as moderately salt sensitive (Mass & Hoffman,

1977). Salinity, as in other plants, has shown to decrease growth, photosynthesis and chlorophyll in lettuce (Han & Lee, 2005; Yan-fang & Jun-yu, 2008).

2.2 Silicon and salinity stress mitigation

Si, the second most abundant element on earth, second only to oxygen (Marshak, 2005:43), is ever-present in plants (Epstein & Bloom, 2005:227-230), in quantities equal or more than many macro nutrients (Epstein, 1999). Si is found as silicic acid (H_4SiO_4) in the soil solution (Epstein, 1999), and when absorbed, it is transported to shoots and polymerises into solid amorphous silica ($SiO_2 \cdot nH_2O$), called opal, found mainly in cell walls (Epstein & Bloom, 2005).

The benefits of Si range from enhancement of plant growth and mechanical strength to improvement of mineral nutrition and resistance to abiotic and biotic stresses (Epstein, 1994). Despite this, Si does not qualify as essential under the widely accepted (yet disputed) definition of plant nutrient essentiality by Arnon & Stout (1939). Epstein (1999), rejects this definition and argues that Si qualifies as quasi-essential, because many plants grown with a deficiency in Si show irregularities in their growth, development or reproduction when compared to plants grown with the element in abundance.


There is a profusion of literature evaluating the benefits of Si. Included in this, is its well documented ability to mitigate stresses associated with salinity in plants, specifically, stresses associated with high levels of NaCl. This is evident in a wide range of plant species, including, but not limited to, wheat (Ahmad, 1992) barley (Liang, 1999), tomato (Zhu, 2004a) cucumber (Zhu, 2004b), zucchini (Savvas, 2009) grapevine (Soylemezoglu, 2009) and sugarcane (Ashraf, 2010).

Mechanisms of mitigation are at variance in the literature, with the majority of studies concluding that Si reduces the uptake, accumulation or concentration of Na in plant tissues (Ahmad *et al.*, 1992, Savvas *et al.*, 2009, Soylemezoglu *et al.*, 2009 and Hashemi, *et al.* 2010). Romero-Aranda *et al.* (2006), conclude that the mitigating properties are due to a Si induced increase in water storage capacity, thus diluting Na in plant tissues. Ma *et al.* (2001:31), conclude that silicon's beneficial affects stem from the decreased Na influx through a Si induced decrease in transpiration.

2.3 Antioxidants and salinity stress

ROS are highly reactive chemical species that, although have recently been shown to have beneficial signalling properties, can damage, impair the function of cells or even cause cell death. They are ubiquitous species in plants and are ever present in cells in aerobic conditions, where O₂ is utilised for energy production (Mittler, 2002; Koolman & Rohm, 2005) as all plant do (Halliwell & Gutteridge, 1999). Examples of ROS include superoxide (O₂⁻), Hydrogen peroxide (H₂O₂) and the hydroxyl radical (OH[•]) (Murray et al., 2003). Due to the high levels of molecular oxygen in the early atmosphere, leading to the inevitable formation of ROS, living organisms have had to evolve mechanisms to counteract the deleterious effects of ROS billions of years ago, in the form of antioxidants (Ndhala et al., 2010). In addition to standard levels of ROS, abiotic stresses, including salinity stress, together with salinity induced osmotic stress, is known to increase the concentrations of ROS (Dat et al., 2000; Xiong & Zhu, 2002).

Antioxidants can simply be defined as any substance that hinders or prevents oxidation of an oxidizable substrate, such as a protein or lipid (Halliwell & Gutteridge, 1999). There are various different antioxidants found in plants, falling into one of two categories; i.e. enzymatic or non-enzymatic (Ndhala et al., 2010). CAT, an indispensable enzymatic antioxidant, and a major antioxidant in plants (Mittler, 2002) is responsible for the conversion of the free radical hydrogen peroxide which is found in high concentrations in chloroplasts and leaf peroxisomes. CAT which is also found predominantly in peroxisomes, understandable due to the high concentrations of H₂O₂, converts H₂O₂ to water and oxygen, expressed by the following equation (Mittler, 2002; Feierabend, 2005:101):



In terms of salinity stress, CAT has shown to decrease in tomato (Zhu et al., 2004a), cucumber (Zhu et al., 2005b), *Plantago media* (Sekmen et al., 2007) both increase and decrease in two different studies on lettuce (Eraslan et al., 2007; Leyva et al., 2011), and increase in *Beta maritima* (Bor et al., 2003) and *Jatropha curcas* (Gao et al., 2008) when plants were exposed to various levels of NaCl. These studies suggest that the effects of NaCl on CAT production varies from one species to the next, however, Mittler (2002) states that abiotic stresses generally lead to the production of CAT.

Phenolics are organic compounds found in all higher plants, comprised of a hydroxyl group bonded to an aromatic hydrocarbon. Gallic acid is a common example, and is used as a representation for total polyphenolic quantifications in some phenolics assays (Figure 2.1).

Polyphenolics on the other hand, are defined by large amounts of these phenol units, and have only comparatively recently been recognized as antioxidants, with their antioxidant properties due to their electron donating activities (Grace, 2005:151-153). They have the ability to scavenge ROS directly due to being less reactive than oxygen, and therefore do not promote further redox reactions.

TP in plants subjected to NaCl salinity stress show a clear tendency to increase over time. Parida et al. (2002) and Parida et al. (2004), studying the effects of NaCl on *Bruguiera parviflora* and *Aegiceras corniculatum*, species of mangrove, reported significant increases in polyphenol concentrations over time. Mangroves are plants that inhabit saline conditions, and consequently have various mechanisms for salt tolerance. An increase in TP concentration with NaCl stress over time in mangroves is therefore a good indication that polyphenolics play a vital role in salinity stress mitigation. Increases in TP with salinity stress is not only reported in the salt tolerant mangroves, but also reported in important agricultural crops, such as maize. Hichem et al. (2009), report vast increases in TP concentrations with increasing NaCl salinity, and conclude that TP play an important physiological role in maize salinity tolerance.

GSH, a molecule with various functions, and like CAT, is a widely known antioxidant (Tausz et al., 2004). GSH reacts with many different ROS, and it plays an essential role in the ascorbate-glutathione cycle (Foyer et al. 2005:1-2). The ascorbate-glutathione cycle results in the naturalisation of H_2O_2 , whereby APX reduces H_2O_2 to water, using ascorbate (vitamin C). GSH plays a later role in this pathway by being oxidised to GSSG, allowing the completion of the cycle with the reduction of dehydroascorbate to ascorbate (Smirnoff, 2005:65).

As stated previously, salinity stress results in increased concentrations of ROS, with an increase in GSH known to infer protection against ROS (Tausz et al., 2004). It is therefore no surprise that salinity stress is widely reported to result in an increase in GSH concentration (Saqib et al., 2008; Leyva et al. 2011), an appropriate response to the salinity induced increase in ROS. Furthermore, Tausz et al. (2004) stated that in addition to the salinity induced increase in GSH concentration, there is a prior increase in the GSH:GSSG ratio, which could act as signals allowing for appropriate cellular reactions (Foyer et al., 2005:1).

2.4 Antioxidant interactions with silicon and salinity stress

As previously discussed, Si has the ability to mitigate the negative effects associated with NaCl induced salinity stress. In addition to the NaCl related amelioration of salinity stress, which is commonly attributed to silicon's influence on Na, studies have indicated that Si can influence antioxidant activity. The effects of Si on antioxidants have been the focus of only a handful of studies. Important crop species that have been studied include; barley (Liang, 1999; Liang et al., 2003), tomato (Zhu, 2004a), cucumber (Zhu, 2004b), maize (Moussa, 2006) spinach (Eraslan *et al.*, 2008) wheat (Saqib, 2008) and alfalfa (Wang et al., 2011).

Investigations in almost all of the studies considering the effects of Si on CAT concentrations in salinity stressed plants, lead to an increase in CAT (Liang et al. 2003; Moussa 2006; Eraslan et al, 2008; Wang et al. 2011), with Zhu et al., (2004a), concluding that Si counteracted the deleterious effects of NaCl by means of increases in CAT. Zhu et al., (2004b), however, found a general decrease in CAT concentrations when investigating cucumber. Gong *et al.* (2005), investigating applications of Si to ameliorate drought stress, a primary limitation for plants growing in saline conditions in the form of an increase in the soil solution's osmotic potential (Marschner, 1995:662-663), found that applications of Si lead to significantly decreased levels of CAT.

Liang et al., (2003), and Saqib et al., (2008), are two of the studies that examined the influence of GSH on salinity toxicity on barley and wheat respectively, as affected by Si applications. Both found that GSH concentrations increased over time compared to NaCl treatments alone. A higher concentration of GSH is advantageous and an appropriate response with regards to stress as additional GSH would result in superior antioxidative defence (Tausz et al., 2004). It follows that a better defence against ROS would result in a decrease in salinity induced production of ROS, and therefore the alleviation of salinity stress (Foyer et al., 2005:1-2,16-18).

Lipid peroxidation is a chain reaction that can lead to continuous tissue injury. This process provides a continuous supply of free radicals, resulting in recurring radical damage (Figure 2.3). To control lipid peroxidation, plants use antioxidants and therefore prevent the otherwise continual tissue damage from this reaction. MDA, a compound resulting only from the degradation of lipids with three or more double bonds, is used as a marker of lipid peroxidation (Murray et al., 2003).

MDA concentrations in NaCl stressed plants show significant increases compared to those without NaCl stress (Liang 1999; Zhu et al., 2004a; Moussa 2006). As could be

hypothesised, applications of Si on salt stressed plants results in a decrease in MDA content, reflecting a decrease in lipid peroxidation and therefore an alleviation of salinity toxicity (Liang 1999; Liang et al., 2003; Zhu et al., 2004; Moussa 2006).

Gathering and considering the information discussed in this literature review, the subject of this thesis was lettuce, one of the most important vegetable crops (Hemy, 1984:104) yet a plant species that to the author's knowledge has not been studied in relation to Si and NaCl salinity stress. The chapters that follow are the results of these investigations.




Figure 2.1: Gallic Acid




Figure 2.2: The ascorbate-glutathione cycle, demonstrating the role of GSH in this pathway. and 4mM Si (Adapted and simplified from Smirnoff, 2005).




Figure 2.3: Lipid peroxidation. In this example, the reaction is initiated by a free radical (X). The process results in Malondialdehyde (MDA) (Adapted from Murray *et al.*, 2003).

CHAPTER THREE

THE ALLEVIATION OF SALINITY INDUCED STRESS WITH THE APPLICATION OF SILICON IN SOILLESS GROWN *Lactuca sativa* L. 'Eish!'.

3.1 ABSTRACT

The effects of 30 and 60 mM NaCl on Lettuce (*Lactuca sativa* L. 'Eish!'), grown in soilless culture, with additions of 0, 1, 2 and 4 mM Si was evaluated. Height, leaf number, weight, chlorophyll content and elemental analysis of plants were examined. Fresh root and shoot weight significantly increased with additions of 2 mM Si compared to that of the 60 mM NaCl control, with notable increases with Si treatments of 1 and 4 mM. Dry root and shoot weight with additions of 1 mM Si, and in shoot weight with additions of 2 mM Si, significantly increased when compared to the 60 mM NaCl control. Increases in dry root and shoot weight of the other Si additions lead to clear, yet insignificant increases in dry weights. Few differences in Cl content were evident amongst treatments, besides a significant increase in root content and a significant decrease in shoot content with applications of 4 mM Si exposed to 60 mM NaCl. Na shoot content of 1, 2 and 4 mM Si at 60 mM NaCl showed a significant decrease compared to that of the control. Applications of 2 and 4 mM Si showed increased shoot Si when compared to both the 30 and 60 mM NaCl controls. With clear evidence for a reduction in Na content in shoots with applications of Si at higher NaCl concentrations, and further increases in plant weights, it can be stated that applications of Si are of a distinct benefit for the growth of lettuce.

Key Words: Lettuce, Si, NaCl, Salt, Sodium chloride, Hydroponics

3.2 INTRODUCTION

Silicon [Si], an element whose abundance in the earth's crust is second only to oxygen (Marshak, 2005:43), can contribute as much as 0.1 - 10% of the dry matter of plants. These levels are equal to or exceeding those of essential macro nutrients (Epstein & Bloom, 2005:52). Si has in the past been classified as a non-essential element for plant nutrition, that is, one which does not meet the criteria for essentiality in the classic definition of essentiality proposed by Arnon & Stout (1939). More recently Si has been

classified as either a beneficial element (Marschner, 1995:417), or as a quasi-essential element in plants (Epstein, 1999).

Contributing to its classification as a beneficial or quasi-essential element in plants is its role as a plant stress ameliorator (Epstein, 1994; Belanger et al., 1995; Epstein, 1999). An important aspect of this is the ability of Si to ameliorate salt toxicity, defined as conditions of high salt build-up, with sodium chloride [NaCl] being the most common cause of saline conditions (Harris, 1992; Epstein & Bloom, 2005:332-334).

Salinity has two methods of causing abnormalities in plant growth. The first, which is independent of the type of salt, is by creating a higher osmotic potential in the medium surrounding the roots, and because osmosis is the principle method of water absorption in plants (Stern, 2006:151-152), it results in a decrease in plant-water availability (Carter, 1981). The second, which is salt dependent, is caused by an excess of available ions. In the case of NaCl salinity, the ions in question are Sodium [Na⁺] and Chloride [Cl⁻]. Cl is known to be an essential micronutrient in plants (Epstein & Bloom, 2005:57; Stern, 2006:161), and some argued that Cl should be classified as a macronutrient, contributing as much as 1% of the dry weight of some plants (Muckle, 1993). Na, on the other hand, is required by only a few plants (Epstein & Bloom, 2005:231-232), and has shown to be the primary cause of growth suppression in lettuce (Tas et al., 2005). In general, high concentrations of both Na and Cl are associated with leaf scorching (Shannon & Grieve, 1999), necrosis and defoliation (Maas & Hoffman, 1977).

The mitigation of NaCl induced toxicity with the application of Si has been demonstrated in many important agricultural and ornamental crops, including Alfalfa (Wang et al., 2011), Tomato (Romero-Aranda et al., 2006), Wheat (Ahmad *et al.*, 1992), Zucchini (Savvas et al., 2009), Grapevine (Soylemezoglu et al., 2009), Mesquite (Bradbury & Ahmad, 1990), Roses (Savvas et al., 2007; Reezi et al., 2009) and several others. The present study investigated the ability of Si to alleviate NaCl toxicity in Lettuce (*Lactuca sativa* L. 'Eish!').

3.3 MATERIALS AND METHODS

The experiment took place within the experimental greenhouse located at the Cape Peninsula University of Technology in Cape Town, South Africa - 33° 55' 58.27''S, 18° 25' 57.04 E. The temperature within the greenhouse ranged from 9°C to 29°C, the mean being 17.6°C and the humidity ranged from 44% to 99%, the mean being 77%.

Lactuca sativa L. 'Eish!' seeds (Hygrotech, South Africa) were sown in vermiculite, and germinated under 40% shade-cloth. Once all seeds had germinated (8 days after sowing) they were fertilised daily with half strength CHEMICULT (Starke Ayres, South Africa) nutrient solution, and increased gradually to full strength (Harris, 1992:55-62). Once the seedlings were fully established (40 days after sowing) 80 plants were randomly selected and transplanted into individual 12.5 cm pots, containing expanded clay as the medium, and placed within the soilless growing system. The sub-irrigation, closed, soilless growing system consisted of a nutrient solution holding tray, delivering a constant ± 1 cm of circulating nutrient solution to each treatment. Using a randomised block design, 10 plants were placed within eight individual soilless culture units, allowing for 10 repeats of each.

3.3.1 Basal Nutrient Solution

Analytical grade, Potassium nitrate [KNO_3], Calcium nitrate [$(\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O})$], Ammonium dihydrogen phosphate [$\text{NH}_4\text{H}_2\text{PO}_4$], and Magnesium sulphate [$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$], were used to add 16 mM N, 6 mM K, 4 mM Ca, 2 mM P, 1 mM S and 1 mM Mg to the nutrient solution, which was based on the modified Hoagland solution presented by Epstein & Bloom (2005:31). Micro nutrients were supplied by using HYGROPLEX (Hygrotech, South Africa), adding 58 μM Boron [B], 11.4 μM Manganese [Mn], 7.6 μM Zinc [Zn], 1 μM Copper [Cu], 0.65 μM Molybdenum [Mo] and 31.9 μM Iron [Fe] to the nutrient solution. The complete basal nutrient solution was prepared using deionised water and had an electrical conductivity (EC) of 1.9 mS cm^{-1} .

3.3.2 Silicon and NaCl Treatments

Four levels of Si, 0 mM, 1 mM, 2 mM and 4 mM, were supplied by adding AGRISIL K50 (PQ Silicas, South Africa) - Potassium silicate [K_2SiO_3], each of which were combined with two levels of NaCl, 30 mM and 60 mM, making a total of 8 treatments. Treatments, which were applied in a single dose, commenced one week after transplantation. Newly prepared nutrient solutions containing 30 mM NaCl had a mean EC of 5.20 mS cm^{-1} and solutions containing 60 mM NaCl had a mean EC of 8.10 mS cm^{-1} . These EC levels were maintained daily. Potassium [K] was balanced in all treatments by subtracting additional K from KNO_3 , and supplementing lost Nitrate [NO_3] by adding Nitric Acid [HNO_3]. The nutrient solution, which was replaced weekly, was maintained daily at a pH of 6.0 using Hydrochloric acid [HCl] and Sodium hydroxide [NaOH].

3.3.3 Plant Growth Parameters

Plant growth, in terms of plant height and leaf number, of all treatments, were measured at weekly intervals. Plant height was measured with a measuring tape, from the surface of the medium to the tip of the tallest leaf.

Five plants per treatment were harvested after 11 weeks from sowing (Hadfield, 2001:110), and separated into roots and shoots. Roots were rinsed once with tap water and twice with deionised water before being patted dry with paper towel. After the fresh weight had been recorded, plants were immediately placed within a specimen oven at 60°C until constant weight, and the dry weight recorded.

3.3.4 Determination of chlorophyll content in plant leaves

Extraction of chlorophyll by Dimethyl sulphoxide [DMSO] was done following a modified method described by Hiscox and Israelstam (1979). This method allows for the extraction of chlorophyll without plant tissue maceration. Five lettuce plants per treatment were lyophilised and ground to a fine powder using a mortar and pestle and stored at -80°C until analysed. 100 mg of the powder was placed in a 15 mL vial containing 7 mL DMSO and incubated at 65°C for 20 minutes. After the incubation period, an additional 3 mL of DMSO was added to the vial, and stored at 4°C. After a 24 hour storage period, the extract was centrifuged at 3220 g for 10 minutes, after which the supernatant was transferred to a new 15 mL vial. Before spectrophotometrical analysis, the supernatant was diluted to a factor of 15 of which 1 mL was placed in a cuvette and absorbance values of 645 and 663 nm recorded. Absorbance values were used in the equation proposed by Arnon (1949), to determine total leaf chlorophyll content against DMSO blank, expressed as mg g⁻¹ dry weight.

3.3.5 Elemental analysis

Dried plant material was used to analyse for Na, Cl and Si. Na and Si was analysed by ashing the ground sample and dissolving the ash in HCl. Elemental concentrations were determined using an inductively-coupled plasma (ICP) emission spectrophotometer. Cl was analysed by digesting the dried, ground sample with HNO₃. An excessive quantity of Silver nitrate [AgNO₃] was added, followed by back titration with Potassium thiocyanate

[KSCN]. Elemental concentrations of Na, Cl and Si are expressed as mg g⁻¹ of dried plant material (A. Van Deventer, Bemlab, South Africa, Personal Communication).

3.3.6 Statistical Analysis

Statistical analysis was performed by using two-way independent analysis of variance (ANOVA) followed by the Bonferroni post test, with Si and NaCl concentrations being the factors assessed. Computations were executed with the software program SPSS (Urdan, 2005:120-170).

3.4 RESULTS

Leaf number showed no significant differences between Si treatments, whether compared to the control, containing 0 mM Si, or to each other (data not shown). Plant height showed similar insignificance between treatments and controls, besides the expected general increase in height over the growing period (Figure 3.1). There was a significant interaction between salt concentration and silicon treatments on fresh root weight ($P \leq 0.05$) and fresh shoot weight ($P \leq 0.1$). Main effect analysis showed a significant decrease in fresh root weight and fresh shoot weight ($P \leq 0.1$ and $P \leq 0.05$ respectively) between the two controls (Figure 3.2 and 3.3). Fresh root weight and fresh shoot weight showed no significant differences at Si levels of 1 mM and 4 mM whether grown with 30 mM or 60 mM NaCl, when compared to their respective control. The treatment of 2 mM Si exposed to 60 mM did however show a significant increase in fresh root ($P \leq 0.05$) and shoot weight ($P \leq 0.1$) when compared to the control. Although only 2 mM Si showed a significant effect on fresh root and shoot weight, there was a marked, yet insignificant, increase in fresh root and shoot weight of 4 mM Si at 30 mM NaCl, and 1 mM and 4 mM Si at 60 mM when compared to the control (Figure 3.2 and 3.3).

There was also a significant interaction between salt concentration and silicon treatments on dry root weight ($P \leq 0.1$) and dry shoot weight ($P \leq 0.05$). Main effect analysis showed that dry root weight significantly decreased ($P \leq 0.1$) between the control of 30 mM NaCl and that of 60 mM NaCl. Besides a significant increase ($P \leq 0.1$) in dry root weight between 1mM Si exposed to 60 mM NaCl and that of the control, the other treatments showed no significant differences, even though there was again a marked increase in weight, especially when compared to the 60 mM NaCl control (Figure 3.4). Dry shoot

weight showed no significant increase between controls. However, there were significant increases ($P \leq 0.05$) in weight of 1 mM and 2 mM Si when compared to the 60 mM NaCl control. Again, although other treatments resulted in insignificant differences, there was a marked increase in root and shoot dry weight when compared to that of the control (Figure. 3.5).

Although there was a significant interaction between salt concentration and silicon treatments on root ($P \leq 0.001$) and shoot ($P \leq 0.05$) Cl content, main effect analysis showed that Cl concentrations only increased significantly in roots of the 4 mM Si with 60 mM NaCl treated plants and the shoots of 2 mM Si plants. This in turn corresponded to a significant decrease ($P \leq 0.1$) in shoot Cl when compared to the control at 60 mM NaCl (Table 3.1) When comparing Cl content of plants treated with the same level of Si, 2 mM Si at 60 mM NaCl showed a significant decrease in shoot Cl content ($P \leq 0.05$) compared to the plants treated with 30 mM NaCl. Contrarily, plants treated with 4 mM at 60 mM NaCl showed a significant increase in root Cl content ($P \leq 0.01$) when compared to the plants treated with 30 mM NaCl (Table 3.2).

There was a significant interaction between salt concentration and silicon treatments on shoot Na content ($P \leq 0.001$) but not on root Na content. Although no significant differences in root Na content were obtained, there was a marked decrease in Na root concentration at 60 mM NaCl (Figure 3.6). Main effect analysis showed that Na shoot content at 30 mM NaCl mirrored the results of that of the roots, and showed no significant differences between treatments. However, Na shoot content at 60 mM NaCl with Si levels of 1, 2 and 4 mM all resulted in a significant decrease in shoot Na content compared to that of the control (Table 3.1). When comparing Na content of plants treated with the same level of Si, root Na significantly increase between the controls of 30 and 60 mM NaCl ($P \leq 0.001$) and shoot Na significantly increase ($P \leq 0.001$) with 60 mM NaCl when compared to 30 mM NaCl at all Si levels (Table 3.2).

There was a significant interaction between salt concentration and silicon treatments on Si shoot content ($P \leq 0.001$), but not on Si root content. Main effect analysis showed that there was however a significant decrease ($P \leq 0.05$) of root Si between treatments at 30 and 60 mM NaCl receiving 1mM Si. Si shoot concentrations of 2 and 4 mM, at both 30 and 60 mM NaCl showed a significant increase when compared to the control. When the two NaCl levels at the same 4 mM Si treatment were compared, a significant decrease ($P \leq 0.01$) in Si was observed in the shoots. This was apparent in the 4 mM treatment alone (Table 3.2).

There was a significant interaction between salt concentration and silicon treatments on chlorophyll content ($P \leq 0.05$). Main effect analysis revealed that, when comparing Si treated plants to the controls, there was only a significant decrease in chlorophyll with applications of 2 mM Si at 30 mM to that of the control (Figure 3.7).

3.5 DISCUSSION

The results obtained from this study support the fundamental findings of previous studies, that Si has beneficial mitigating properties with regards to NaCl induced salinity stress. These findings indicate some benefits of Si applications at 30 mM NaCl, however, the benefits of Si seem to be more apparent with the higher experimental concentrations of 60 mM NaCl. This is evident when looking at the significant increase in fresh root and shoot weight at 2 mM Si at 60 mM NaCl, the significant increases in dry root, at 1 mM Si, dry shoot weight at 1 and 2 mM, and the overall marked increases in fresh and dry weights when comparing all Si treatments to that of the 60 mM NaCl control (Figure 3.2, 3.3, 3.4 and 3.5). These findings are supported by the literature which states that dry shoot and root weights of plants increase with applications of Si compared to a range of NaCl concentrations lacking Si (Romero-Aranda et al., 2005; Tuna et al., 2008; Ashraf et al., 2010).

Chlorophyll content showed no significant differences, with only one exception with an application 2 mM Si at 30 mM NaCl (Figure 3.7). This is in line with Tas et al. (2005), who although did not investigate Si, reported negligible effects of NaCl on chlorophyll content on lettuce. However, in a study conducted by Hashemi et al. (2010), evaluating the mitigating effects of Si on Canola, a vast increase in chlorophyll content was found.

A notable oddity with regards to elemental analysis was the exceptional accumulation of Cl in the roots of the plants treated with 4 mM Si at 60 mM (Table 3.1 and 3.2). This, when compared to the control, corresponded to a significant decrease in Cl accumulation in shoots. Savvas et al. (2007), and Savvas et al. (2009), found similar results, reporting a significant decrease in shoot Cl content with Si applications. However, Romero-Aranda et al. (2005), and Soylemezoglu et al. (2009), reported a slight increase in Cl shoots concentration with applications of Si.

A significant finding with regards Na content was the reduction of Na within shoots when Si treated plants exposed to 60 mM NaCl when compared to the control (Table 3.1). The

decrease of Na content in Si treated plants has been a major finding when evaluating the ability of Si to mitigate NaCl salinity stress. Many studies that evaluate Si and NaCl relate NaCl mitigation to a reduction in Na shoot concentrations (Saquib et al., 2008; Tuna et al., 2008; Savvas et al., 2009; Ashraf *et al.* 2010), the reduction of which may be linked to Si reducing transpiration, as concluded by Ma et al. (2001). Savvas (2007), also reported a decrease in Na with applications of Si, but did not conclude this to be factor influencing the ability of Si to mitigate salinity stress.

Other mechanisms of salinity mitigation by Si have been suggested, including the ability of Si to improve water storage within plant tissues. Due to the higher growth rate that this would allow, it follows that salt concentrations would decrease through dilution, and therefore ameliorate salt toxicity (Romero-Aranda et al., 2005). Anti oxidants have also been a topic of investigation, with Liang (1999), Zhu et al. (2004a) and Zhu et al. (2004b), investigating the interaction of Si with antioxidants in salt stressed barley, cucumber and tomato respectively, with promising results.

Although transpiration and anti oxidants were not factors investigated in this study, plant weight increases, and subsequent decreases in Na shoot concentrations, leads to the conclusion that mitigation of NaCl induced salinity with applications of Si was at least partially due to the reduction of shoot Na content. Additions of Si to plants under salinity stress can be recommended, especially at higher salinity levels, as shown by the results of this paper.

3.6 ACKNOWLEDGMENTS

This study was funded by the Cape Peninsula University of Technology, and supported by the Mauerberger Foundation Scholarship.

.

3.7. REFERENCES

Ahmad, R., Zaheer, S.H. & Ismail, S. 1992. Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Science*, 85(1):43-50.

Arnon, D.L. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidases in *Beta vulgaris*. *Plant Physiology*. 24:1-15.

Arnon, D.L. & Stout, P.R. 1939. The essentiality of certain elements in minute quantities with special reference to copper. *Plant Physiology*, 14: 371-375.

Ashraf, M., Rahmatullah., Afzal, M., Ahmed, R., Mujeeb, F., Sarwar, A. & Ali, L. 2010. Alleviation of detrimental effects of NaCl by silicon nutrition in salt-sensitive and salt-tolerant genotypes of sugarcane (*Saccharum officinarum* L.). *Plant and Soil*, 326(1):381-391.

Belanger, R.R, Bowen, P.A., Ehret, D.L. & Menzies, J.G. 1995. Soluble silicon: it's role in crop and disease management of greenhouse crops. *Plant Disease*, 79(4):329-336.

Bradbury, M. & Ahmad, R. 1990. The effect of silicon on the growth of *Prosopis juliflora* growing in saline soil. *Plant and Soil*, 125:71-74.

Carter, D.L. 1981. Salinity and plant productivity. *CRC Handbook of Agricultural Productivity*, 1:117-133.

Epstein, E. 1994. The anomaly of silicon in plant biology. *Proceedings of the National Academy of Sciences of the United States of America*, 91:11-17.

Epstein, E. 1999. Silicon. *Annual Review of Plant Physiology and Plant Molecular Biology*, 50: 641-664.

Epstein, E. & Bloom, A. J. 2005. *Mineral nutrition of plants: Principles and perspectives*. 2nd Ed. Sunderland, MA: Sinauer Associates.

Hadfield, J. 2001. *The A-Z of vegetable gardening in South Africa*. Cape Town: Struik.

Harris, D. 1992. *Hydroponics: The complete guide to gardening without soil*. Cape Town: New Holland Publishing.

Hiscox, J.D. & Israelstam, G.F. 1979. A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany*, 57:1332-1334.

Hashemi, A., Abdolzadeh, A. & Sadeghipour, H.R. 2010. Beneficial effects of silicon nutrition in alleviating salinity stress in hydroponically grown canola, *Brassica napus* L., plants. *Soil Science and Plant Nutrition*, 56(2):244-253.

Liang, T. 1999. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. *Plant and Soil*, 209:217-224.

Maas, E.V. & Hoffman, G.J. 1977 . Crop salt tolerance – Current Assesment. *Journal of Irrigation and Drainage Division*, 103:115-134.

Ma, J.F., Miyake, Y. & Takahashi, E. 2001. Silicon as a beneficial element for crop plants. In Datnoff, L.E., Snyder, G.H. & Korndörfer, G.H. (Eds). *Silicon in agriculture*. New York: Elsevier.

Marshak, S. 2005. *Earth: Portrait of a Planet*. 2nd Ed. New York: W.W. Norton and Company.

Marschner, H. 1995 . *Mineral Nutrition of Higher Plants*. 2nd Ed. London: Academic Press.

Muckle, M.E. 1993. *Hydroponic nutrients: Easy ways to make your own*. Canada: Growers Press.

Reezi, S., Babalar, M. & Kalantari, S. 2009. Silicon alleviates salt stress, decreases malondialdehyde content and affects petal color of salt stressed cut rose (*Rosa xhybrida* L.) 'Hot Lady'. *African Journal of Biotechnology*, 8(8):847-855.

Romero-Aranda, M.R., Jurado, O. & Cuartero, J. 2006. Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *Journal of Plant Physiology*, 163(8):847-855.

Savvas, D., Giotis, D., Chatzieustratiou, D., Bakea, M. & Patakioutas, G. 2009. Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environmental and Experimental Botany*, 65(1):11-17.

Savvas, D., Kizas, G., Lydakis-Simantiris, N., Salahas, G., Papadimitriou, M. & Tsouka N. 2007. Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *European Journal of Horticultural Science*, 72(2):73-79.

Shannon, M.C. & Grieve, C.N. 1999. Tolerance of vegetables to salinity. *Scientia Horticulturae*, 78:5-38.

Saqib, M., Zorb, C. & Schubert, S. 2008. Silicon-mediated improvement in the salt resistance of wheat (*Triticum aestivum*) results from increased sodium exclusion and resistance to oxidative stress. *Functional Plant Biology*, 35(7):633-639.

Soylemezoglu, G., Demir, K., Inal, A. & Gunes, A. 2009. Effect of silicon on antioxidant and stomatal response of two grapevine (*Vitis vinifera* L.) rootstocks grown in boron toxic, saline and boron toxic-saline soil. *Scientia Horticulturae*, 123(2): 240-246.

Stern, K.R. 2006. *Introductory plant biology*. 10th Ed. New York: McGraw Hill.

Tas, G., Papadandonakis, N. & Savvas, D. 2005. Response of lettuce (*Lactuca sativa* L. var. *longifolia*) grown in a closed hydroponic system to NaCl- or CaCl₂-salinity. *Journal of Applied Botany and Food Quality*, 79(2):136-140.

Tuna, A.L., Kaya, C., Higgs, D., Murillo-Amador, B., Aydemir, S. & Girgin, A.R. 2008. Silicon improves salinity tolerance in wheat plants. *Environmental and Experimental Botany*, 62(1):10-16.

Urduan, T.C. 2005. *Statistics in Plain English*. 2nd Ed. New Jersey: Lawrence Erlbaum Associates.

Wang, X., Wei, Z., Liu, D. & Zhao, G. 2011. Effects of NaCl and silicon on activities of antioxidant enzymes in roots, shoots and leaves of alfalfa. *African Journal of Biotechnology*, 10(4):545-549.

Zhu, Z., Wei, G., Li, J., Qian, Q. & Yu, J. 2004a. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Science*, 167: 527–533.

Zhu, Z., Al-aghabary, K. & Shi, Q. 2004b. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *Journal of Plant Nutrition*, 27(12):2101-2115.





Figure 3.1: The effect of 30 and 60 mM NaCl on plant height, with additions of 0,1, 2 and 4 mM Si. Measurements were take weekly from 27 May (week 1) to 17 June (week 4) 2011.






Figure 3.3. The effect of 30 and 60 mM NaCl on fresh shoot weight, with additions of 0, 1, 2 and 4 mM Si.




Figure 3.4. The effect of 30 and 60 mM NaCl on dry root weight, with additions of 0, 1, 2 and 4 mM Si.




Figure 3.5. The effect of 30 and 60 mM NaCl on dry shoot weight, with additions of 0, 1, 2 and 4 mM Si




Figure 3.6. The effect of 30 and 60 mM NaCl on Na root content, with additions of 0, 1, 2 and 4 mM Si




Figure 3.7. The effect of 30 and 60 mM NaCl on total chlorophyll content with additions of 0, 1, 2 and 4 mM Si.

Table 3.1. The effect of 30 and 60 mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4 mM Si.

Treatments		Cl (mg g ⁻¹)		Na (mg g ⁻¹)		Si (mg g ⁻¹)	
NaCl	Si	Roots	Shoots	Roots	Shoots	Roots	Shoots
30mM	0 mM	12.18	16.78	7.17	11.11	0.42	0.06
	1 mM	9.78	18.70	8.12	11.41	0.44	0.12
	2 mM	9.60	22.54	8.59	11.22	0.51	0.15**
	4 mM	14.42	13.74	8.10	11.64	0.43	0.43***
60 mM	0 mM	15.98	21.20	14.30	16.88	0.37	0.07
	1 mM	14.82	13.62	9.43	13.75***	0.29	0.11
	2 mM	7.66	14.20	10.77	15.40**	0.45	0.16***
	4 mM	74.8***	12.74*	8.83	14.00***	0.40	0.18***

Values presented are means \pm SE, n = 5. *, **, *** = significant at $P \leq 0.1^*$, $P \leq 0.05^{**}$ or $P \leq 0.01^{***}$. Significance compared to that of 0 mM Si, with corresponding NaCl concentrations (Controls).

Table 3.2. The effect of 30 and 60 mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4 mM Si.

Treatments		Cl (mg g ⁻¹)		Na (mg g ⁻¹)		Si (mg g ⁻¹)	
Si	NaCl	Roots	Shoots	Roots	Shoots	Roots	Shoots
0mM	30 mM	12.18	16.78	7.17	11.11	0.42	0.06
	60 mM	15.98	21.20	14.30***	16.88***	0.37	0.07
1mM	30 mM	9.78	18.70	8.12	11.41	0.44	0.12
	60 mM	14.82	13.62	9.43	13.75***	0.29**	0.11
2mM	30 mM	9.60	22.54	8.59	11.22	0.51	0.15
	60 mM	7.66	14.20**	10.77	15.40***	0.45	0.16
4mM	30 mM	14.42	13.74	8.10	11.64	0.43	0.43
	60 mM	74.8***	12.74	8.83	14.00***	0.40	0.18***

Values presented are means \pm SE, n = 5. *, **, *** = significant at $P \leq 0.1^*$, $P \leq 0.05^{**}$ or $P \leq 0.01^{***}$. Significance compared to that of the same Si concentrations, with different NaCl concentrations.

CHAPTER FOUR

SALINITY INDUCED CHANGES IN OXIDATIVE STRESS AND ANTIOXIDANT STATUS AS AFFECTED BY APPLICATIONS OF SILICON IN LETTUCE (*Lactuca sativa* L. 'Eish!').

4.1 ABSTRACT

The antioxidant content, capacity and certain oxidative stress parameters were investigated in salinity stressed *Lactuca sativa* L. 'Eish!' with the addition of exogenous silicon (Si). No significant changes were shown for total polyphenol concentrations in any of the experimental plants. Applications of 4mM Si significantly increased oxygen radical absorbance capacity (ORAC) values at both 30 and 60 mM NaCl. The redox status of glutathione was also improved with a significant increase in reduced glutathione (GSH) in 2mM Si treated plants at 30 mM NaCl. Increases in both GSH and oxidised glutathione (GSSG) were noted at 60 mM NaCl. The activity of catalase (CAT) was significantly decreased with the addition of 1 and 4 mM Si at 30 mM NaCl. Lipid peroxidation (measured as thiobarbituric acid reactive substances, TBARS) was not influenced by the different Si concentrations. These results suggest that applications of Si could be beneficial with regards to the modulation of oxidative stress in salinity stressed lettuce, and should be considered as an addition when cultivating lettuce.

Key Words: Potassium silicate, ROS, Hydroponics, Catalase, Lipid peroxidation, Glutathione, Polyphenolics, Oxygen Radical Absorbance Capacity, Thiobarbituric Acid Reactive Substances.

4.2 INTRODUCTION

Salinity is a significant plant stress factor and a major problem on all continents (Ghassemi et al., 1995). Saline conditions influence plant growth in two core ways; by increasing the osmotic potential of the soil solution, which has a direct effect on plant-water availability (Carter, 1981), and through ion toxicities, determined by the specific ions that are in excess (Marschner, 1995:662). In most cases, sodium (Na⁺) and chloride (Cl⁻) ions are the cause of saline conditions (Harris, 1992:215; Epstein & Bloom, 2005:332-

334). The effect of NaCl induced salinity on plants differ from plant to plant, but generally leads to a reduced growth rate, sometimes nutritional deficiencies, and at higher concentrations, the accumulation of Na⁺ and Cl⁻, resulting in leaf scorching, chlorosis and necrosis (Shannon & Grieve, 1999; Wahid et al., 1999). Along with these symptoms, salinity stress is known to result in an excess production of reactive oxygen species (ROS) (Jaspers et al., 2005), oxidative damage and a change in concentrations of antioxidants (Bor et al., 2003; Sekmen et al., 2007; Gao et al., 2008). Consequently, ROS are good cellular indicators of stress (Mittler, 2002).

Silicon (Si), a quasi-essential element for plants (Epstein, 1999), has proved to have significant ameliorative properties regarding NaCl toxicity in many important agricultural and ornamental crops. It has shown to repeatedly result in an increase in yield together with other beneficial growth parameters (Romero-Aranda et al., 2006; Savvas et al., 2007; Soylemezoglu et al., 2009). Silicon allows plants to grow more productively on saline land, leading to definite economic benefits, especially for developing nations (Shay, 1990).

The literature disagrees in the manner by which silicon mitigates NaCl toxicity. Si has the ability to influence Na absorption, distribution or transportation (Ahmad et al., 1992; Savvas et al., 2009; Soylemezoglu et al., 2009; Hashemi, et al. 2010), which is sometimes associated with an increase in potassium (K) concentration (Saqib et al., 2008; Tuna et al., 2008; Ashraf et al., 2010). Ameliorative properties of silicon have also been associated with the decrease in plant Na concentrations through dilution, thus mitigating salt toxicity (Romero-Aranda et al., 2006). Ma et al. (2001), suggested silicon's beneficial effects stems from the decreased Na influx through a Si induced decrease in transpiration. In addition to the Si – Na related amelioration of salinity stress, studies have indicated that Si can influence antioxidant activity. This has been reported in, amongst others, salt stressed barley, (Liang, 1999), cucumber, (Zhu et al., 2004a), tomato (Zhu et al., 2004b) and wheat (Saqib et al., 2008).

In this study the influence of Si in NaCl stressed *Lactuca sativa* L. 'Eish!' was investigated in order to assess whether the known Si induced yield increase can be associated with antioxidant concentrations and oxidative damage. Furthermore, it is to our knowledge that this is the first study examining Si applications influencing antioxidant status and oxidative damage in NaCl salinity stressed lettuce.

4.3 MATERIALS AND METHODS

4.3.1 Plant Conditions and Cultivation

Lettuce cultivation took place in the research greenhouse at the Cape Peninsula University of Technology (Cape Town, South Africa). Greenhouse temperatures ranged between a minimum of 9 and maximum of 29°C, and humidity between a minimum of 44 and a maximum of 99%. Seeds were sown in a 128 plug polystyrene seedling tray, with vermiculite as the sowing medium. Seeds germinated 8 days after sowing, and were fertilised daily with half strength NUTRIFEED (Starke Ayres, South Africa); a complete, water soluble fertiliser providing 9.3 mM N, 1.7 mM P, 6.7 mM K, 3.5 mM Ca, 1.8 mM Mg, 4.7 mM S, 53.7 µM Fe, 8.7 µM Mn, 44.4 µM B, 1.5 µM Zn, 6.3 µM Cu and 0.2 µM Mo. Four weeks after sowing, 80 seedlings were chosen at random, and transplanted into the soilless growing systems. Seedlings were transplanted into 12.5cm plastic pots with HYDROTON (Germany) expanded clay as the medium, and divided evenly amongst the 8 soilless growing systems, equating to 8 different treatments with 10 repeats of each. Each closed, sub irrigative, soilless growing system was comprised of a nutrient solution holding tray, delivering ±1cm of recirculating nutrient solution.

4.3.2 Basal Nutrient Solution

Salts used to add macro nutrients to the nutrient solution were of analytical grade; KNO_3 , $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, $\text{NH}_4\text{H}_2\text{PO}_4$, and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ were used to include 16 mM N, 6 mM K, 4 mM Ca, 2 mM P, 1 mM S and 1 mM Mg respectively, based on the modified Hoagland solution as described by Epstein and Bloom (2005:31). Micro nutrients were supplied using HYGROPLEX (Hygrotech, South Africa); a complete commercially available micro nutrient mix, adding 58 µM B, 11.4 µM Mn, 7.6 µM Zn, 1 µM Cu, 0.65 µM Mo and 31.9 µM Fe to the nutrient solution. The basal nutrient solution, having had an electrical conductivity (EC) of 1.9 mS cm^{-1} , was formulated using deionised water.

4.3.3 Silicon and NaCl Treatments

Silicon was added to the basal nutrient solution using K_2SiO_3 (AGRISIL K50), at concentrations of 0 mM, 1 mM, 2 mM and 4 mM Si. Two concentrations of NaCl, 30mM and 60 mM, were applied to each of the 4 Si concentrations, making a total of 8

treatments. Treatments were applied in a single dose, implemented one week after transplantation. Newly prepared treatments containing 0, 1, 2 and 4 mM Si and 30 mM NaCl had a mean EC of 5.20 mS cm⁻¹, and treatments containing 0, 1, 2 and 4 mM Si and 60 mM NaCl, had a mean EC of 8.10 mS cm⁻¹. These EC levels were maintained daily. K that was added with the applications of K₂SiO₃, was balanced in all treatments by subtracting additional K from KNO₃, and supplementing lost NO₃ by adding HNO₃. The nutrient solution, which was renewed weekly, was maintained daily at a pH of 6.0 using HCl and NaOH.

4.3.4 Sampling and Harvesting

4.3.4.1 Antioxidant status and oxidative damage

The shoots of 5 plants per treatment were harvested 6 weeks after being transplanted into the soilless growing system. Once harvested, shoots were rinsed twice in deionised water and patted dry with paper towel. The shoots were then placed directly in an air tight plastic bag, and stored at -80°C. The shoots were transported on dry ice, and lyophilised at -80°C for 16 hours. Once the lyophilisation was complete, samples were ground to a powder using a mortar and pestle and stored at -80°C until analysed.

4.3.4.2 Plant yield and elemental analysis

Plant yield is expressed as fresh and dry weight of roots and shoot in grams. Five plants per treatments were separated into roots and shoots, rinsed twice in deionised water and patted dry with paper towel. After recording the fresh weights, the plant material were placed into paper bags and dried at 65°C. Once the weight of the dried plant material stayed constant, dry shoot and root weights were recorded.

Quantities of shoot Si of the 5 treatments were measured by dissolving the ground, ashed material in HCl. Concentrations, determined using an inductively-coupled plasma emission spectrophotometer, are expressed in mg g⁻¹ of dried plant material.

4.3.5 Total protein analysis

For total protein extraction, 125 mg of lyophilised plant material was homogenised with 6mL of 25mM HEPES-KOH buffer containing 0.2mM EDTA and 2% PVP (pH 7.8), on ice. The homogenate was centrifuged in 2 mL microcentrifuge tubes at 15 000 *g* for 10 minutes at 4°C. The resulting supernatant was transferred to new 2mL microcentrifuge tubes and stored at -80°C until needed.

Protein content was determined using the PIERCE BCA protein assay kit, utilising the procedures accompanying the kit, with absorbance levels being measured at 562 nm. Total protein content (expressed as $\mu\text{g mL}^{-1}$) was used for catalase activity quantification.

4.3.6 Polyphenol antioxidant content and capacity

4.3.6.1 Total polyphenol analysis

For TP analysis, 125 mg of lyophilised plant material was homogenised with 6mL of methanol on ice, and centrifuged in 2mL microcentrifuge tubes at 15 000 *g* for 10 minutes at 4°C. 25 μL undiluted supernatant was then transferred, in triplicate, to a 96 well plate containing 125 μL of 10 times diluted Folin-Ciocalteu's phenol reagent (0.2 N). After 5 minutes, 100 μL of sodium carbonate (7.5%, w/v) was added and the resulting mixture incubated at room temperature for a further two hours. Absorbance values at 765 nm were recorded and compared to those of a Gallic acid standard. Results are expressed as mg GAE g^{-1} .

4.3.6.2 Oxygen radical absorbance capacity analysis

Extraction followed that of TP, with 12 μL of 10 times diluted supernatant being added to 138 μL fluorescein and 50 μL AAPH, and analysed, in triplicate, using a fluorescence spectrophotometer until zero fluorescence occurred. The ORAC values were calculated by dividing the sample curve by that of the Trolox standard, and expressed as $\mu\text{mol TE g}^{-1}$ (Ou et al., 2001).

4.2.7 Catalase analysis

Catalase (CAT) activity was measured, in triplicate, by placing 20 μL of the supernatant (obtained from protein analysis, refer 2.5) with 170 μL of a 50 mM Potassium phosphate

buffer (pH7.0) and 75 μL H_2O_2 in a 96 well micro plate. The disappearance of H_2O_2 was measured over a 2 minute period by measuring absorbance levels at 240 nm every 15 seconds. Catalase activity was calculated using the extinction coefficient of H_2O_2 and expressed as $\mu\text{mol min}^{-1} \mu\text{g}^{-1}$ protein (Aebi, 1984).

4.3.8 Glutathione analysis

For GSH extraction, 125 mg of lyophilised plant material was homogenised with 6 mL of 5% Trichloroacetic Acid buffer, on ice. The homogenate was centrifuged in 2 mL microcentrifuge tubes at 15 000 g for 10 minutes at 4°C . The resulting supernatant was transferred to new 2 mL microcentrifuge tubes and stored at -80°C until needed.

Glutathione concentrations were measured, in triplicate, by placing 50 μL of 50 times diluted supernatant, with 50 μL of 0.3 mM DTNB (5,5'-Dithiobis(2-nitrobenzoic acid)) and 50 μL Glutathione reductase (GR) solution (0.02U/ μL) in a 96 well micro plate. After a 5 minute incubation period at room temperature, 50 μL of 1 mM NADPH was placed in each well. Absorbance levels at 412 nm were measured at 15 second intervals for 5 minutes. Absorbance levels were compared to those of a GSH standard curve, and total GSH expressed as $\mu\text{mol g}^{-1}$ dried plant material.

GSSG extraction followed that of GSH, except for the addition of 4.8 mg of M2VP (1-Methyl-2-vinylpyridinium) within the homogenate buffer. Measurement of GSSG followed that of GSH, except a 25 times diluted supernatant was used, and absorbance levels were compared to those of a GSSG standard curve (Asensi et al., 1999).

4.3.9 Lipid peroxidation analysis

For thiobarbituric acid reactive substances extraction, the method by Yagi, 1984, adapted for plant samples, was utilised. 125 mg of lyophilised plant material was homogenised with 6mL of Methanol, on ice. The homogenate was centrifuged in 2 mL microcentrifuge tubes at 15 000 g for 10 minutes at room temperature. The resulting supernatant was transferred to new 2 mL microcentrifuge tubes and stored at -80°C until needed.

250 μL of undiluted supernatant, with 375 μL of 0.44 mM PCA (perchloric acid) and 125 μL of 42 mM TBA (2-Thiobarbituric acid) were placed in 2 mL microcentrifuge tubes. After a 60 minute incubation period at 100°C, the 2 mL microcentrifuge tubes were centrifuged at 10 000 g for 5 minutes at room temperature. The resulting supernatant was placed, in triplicate, in a 96 well plate, and absorbance levels at 532 nm were compared to those of the malondialdehyde standard and expressed as $\mu\text{mol g}^{-1}$ dried plant material (Yagi, 1984).

4.3.10 Statistical Analysis

Statistical analysis was performed using two-way analysis of variance (ANOVA), followed by the Bonferroni post test, with P values ≤ 0.05 considered significant. Pearson correlations coefficient was used to calculate correlations. Computations were executed with the software program SPSS (Urdu, 2005:120-170).

4.4 RESULTS

Plant weights showed no significant differences when treated plants were compared to the 30 mM NaCl control. However, there were a 12 and 17% increase in fresh shoot and root weight and a 46% increase in dry shoot weight when comparing the 4 mM Si treatment to the control (Table 4.1).

However, plant weights showed a clear increase when treated plants were compared to the 60 mM NaCl control. Although 2 mM Si was the only treatment that resulted in a marginally significant increase ($P \leq 0.1$) in fresh shoot weight, the increase was a considerable (72%). The fresh root weight of the 2 mM Si treated plants resulted in a significant increase (75%) when compared to the control. Dry shoot weights significantly increase by 74 and 75% with both 1 and 2 mM Si treated plants respectively when compared to the control. Although the 4 mM Si treated plants showed no significant difference when compared to the 60 mM NaCl control, there was a marked increase of 56%. Dry root weights showed a marginal significant increase of 68% with 1 mM Si when compared to the 60 mM NaCl control. Again, although dry root weights of 2 and 4 mM Si treated plants resulted in no significant differences when compared to the control, there was a noteworthy increase of 50 and 43% respectively.

Shoot Si concentrations of 1 mM Si treated plants showed no significant difference at both 30 and 60 mM NaCl when compared to their individual controls. Nevertheless, treatments of 2 and 4 mM Si at 30 mM NaCl resulted in a significant concentration increase of 150% and 616% respectively. Similarly, treatments of 2 and 4 mM Si at 60 mM NaCl resulted in a significant increase of 129% and 157% respectively (Figure 4.1.).

With regards to antioxidant status and oxidative stress between the two controls, there were no significant differences between any of the parameters, besides that of the TP controls. The 60 mM NaCl control expressed an 18% increase in TP concentrations when compared to that of the 30 mM NaCl control, resulting in a marginal difference ($P \leq 0.1$) between the two controls. However, when Si treated plants were compared to that of the controls, TP concentrations showed no differences (Table 4.2). ORAC values showed a tendency to increase with applications of Si at all concentrations when compared to the controls. However, only 4 mM Si at both 30 and 60 mM NaCl showed significant increases when compared to controls, with a 28% and 21% increase respectively (Figure 4.2).

Catalase activities showed a clear tendency to decrease with applications of Si when plants were exposed to 30 mM NaCl, with 1 and 4 mM Si, a decrease of 29 and 25% respectively, being significant when compared to the 30 mM NaCl control. Conversely Si applications at 60 mM NaCl showed no tendency to decrease, and showed no significance when compared to that of the 60 mM control (Figure 4.3).

Concentrations of both GSH and GSSG, like that of ORAC, tended to increase with applications of Si. Concentrations of GSH at 2 mM Si with 30 mM NaCl, and both 1 and 4 mM Si with 60 mM NaCl showed significant increases of 138%, 106% and 99% respectively, when compared to their individual controls (Figure 4.4). Concentrations of GSSG showed a significant increase of 100% at 4 mM Si, and a marginally significance increase of 93% ($P \leq 0.1$) at 1 mM Si when compared to the 60 mM NaCl control. The ratio of GSH:GSSG showed a limited trend to increase or decrease, with an application of 2 mM Si with 30 mM NaCl being the only treatment showing a significant increase of 33% ($P \leq 0.1$) when compared to the control.

Like that of TP concentrations, TBARS values showed no significant differences when any of the treatments were compared to their respective controls. In addition to this, no correlations were found when comparing variables within treatments.

4.5 DISCUSSION

Plant stresses, including salinity stress, are known to disturb cellular homeostasis, enhancing the production of ROS (Dat et al., 2000). Additionally, osmotic stress, one of the foremost stresses associated with high salinity levels, has shown to cause the production of ROS (Xiong and Zhu, 2002).

Although ROS have roles as signalling molecules, active generation of which can be initiated by abiotic stresses (Desikan et al., 2005), ROS are traditionally thought of as species causing cell injury or death (Mittler, 2002). In the light of this, plants need ways to detoxify ROS. CAT is a major ROS scavenging enzyme seen as indispensable, with stress bringing about the production of CAT (Mittler, 2002). With ROS causing the production of CAT, it can be argued that lower concentrations of CAT, indicating lower levels of ROS, would be a sign of less oxidative stress. In this study, with the application of Si, there was an overall tendency for CAT concentrations to decrease, significantly with 1 and 4 mM at 30 mM NaCl. Zhu et al. (2004a), however, showed that Si raises CAT levels in salt treated tomatoes, while Zhu et al., (2004b) reported tendencies for CAT levels to decrease with applications of Si on salt treated plants. Bor et al., (2003), Gao et al., (2008) and Eraslan et al., (2007), although not investigating Si, found applications of NaCl to significantly increase CAT activities, supporting the idea that increased CAT activities could be indicative of oxidative stress. Gong et al. (2005) when investigating applications of Si to ameliorate drought stress, found that applications of Si lead to significantly decreased CAT activity. This is relevant to NaCl stress, as osmotic stress is a primary limitation for plants growing in saline conditions (Marschner, 1995:662).

Contrary to CAT activities, yet similarly indicating a decrease in oxidative stress, ORAC levels showed a tendency to increase, significantly at 4 mM Si with both 30 and 60 mM NaCl. ORAC, a hydrophilic antioxidant capacity assay, is one of the most commonly used methods for measuring antioxidative capacity in biology (Yeum et al., 2010). It is assumed that with an increase in TP concentrations, there would be a correlative increase in ORAC values. The level of TP in Si treated plants showed no significant differences between controls, and perhaps even a tendency to decrease. Liu et al. (2007) reported similar results, although with relatively higher TP concentrations, with no correlation to the DPPH assay, a procedure that has shown to be comparable to that of the ORAC assay (Thaipong et al., 2006), even though different mechanisms are involved (Ndhlala et al., 2010).

TBARS concentrations, a well-known indicator of lipid peroxidation (Wada et al., 2008), a process that can lead to cell death (Mittler, 2002), showed no significant differences between the treated and the control plants, and no trend to increase or decrease. This is in disagreement with Liang, (1999), Zhu et al., (2004a), and Zhu et al., (2004b) all finding applications of Si to decrease lipid peroxidation.

Increases in GSH concentrations are known to be associated with salinity stress (Ruiz & Blumwald, 2002; Leyva et al., 2011). The results from this study show that applications of Si tend to increase GSH concentrations, significantly with 2 mM Si at 30 mM NaCl and with 1 and 4 mM Si at 60mM NaCl, when compared to NaCl alone. These results are in agreement with Saquib et al. (2008), reporting that GSH increases with applications of Si in NaCl stressed wheat. Foyer et al., (2005:1) argue that the signal responses that could be brought about by the interaction between GSH and the GSH:GSSG ratio are linked to plant growth cessation. This is interesting when considering that, contrary to conventional thought, active plant growth cessation, as opposed to stress limiting growth, has been linked to surviving adverse environmental conditions (Harberd et al., 2009), including salinity stress (Magome et al., 2008). However, applications of Si are widely reported to increase plant growth parameters (Romero-Aranda et al., 2006; Soylemezoglu et al., 2009), which is in line with the data presented in this study. Therefore, applications of Si induced increases in GSH concentrations seem unlikely to be associated with a growth cessation stress response. GSH are well known antioxidants, and higher concentrations would infer superior antioxidative defence (Tausz et al., 2004), and would therefore logically result in a decrease in ROS concentrations brought about by salinity stress (Foyer et al., 2005:1-2,16-18). This approach would arguably provide more probable grounds for the increases in GSH with applications of Si.

From this study, the overall impression of Si applications in relation to antioxidant concentrations and oxidative stress modulation is that fertilisation and regular inclusion of Si in nutrient solutions, where Si is often limited, is of value to lettuce cultivation. Investigations into the effects of higher salinity concentrations would likely show a clear cut results with regards to the effects of Si applications on antioxidant status in salinity stressed lettuce, and is a recommended focus for future study. Additionally, more insight into the relationships between yield parameters and antioxidant status would bring light to the positive effects Si has on the antioxidative status of plants on a more practical, economically beneficial level.

4.6 ACKNOWLEDGMENTS

This study was funded by the Cape Peninsula University of Technology, and supported by the Mauerberger Foundation Scholarship.

4.6 REFERENCES

- Aebi, H. 1984. Catalase in vitro. *Methods in Enzymology*. 105:121–126.
- Ahmad, R., Zaheer, S.H. & Ismail, S. 1992. Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Science*, 85(1):43-50.
- Asensi, M., Sastre, J., Pallardo, F.V., Lloret, A., Lehner, M., Garcia-de-la Asuncion, J. & Viña J. 1999. Ratio of reduced to oxidised glutathione as indicator of oxidative stress status and DNA damage. *Methods in Enzymology*, 1999(299):267-76.
- Ashraf, M., Rahmatullah., Afzal, M., Ahmed, R., Mujeeb, F., Sarwar, A. & Ali, L. 2010. Alleviation of detrimental effects of NaCl by silicon nutrition in salt-sensitive and salt-tolerant genotypes of sugarcane (*Saccharum officinarum* L.). *Plant and Soil*, 326(1):381-391.
- Bor, M., Özdemir, F. & Türkan, I. 2003. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Science*, 164:77-84.
- Carter, D.L. 1981. Salinity and plant productivity. *CRC Handbook of Agricultural Productivity*, 1:117-133.
- Dat, J., Vandenabeele, S., Vranová, E., Van Montagu, M., Inzé, D. & Van Breusegem, F. 2000. Dual action of the active oxygen species during plant stress responses. *Cellular and Molecular Life Sciences*, 57:779–795.
- Desikan, R., Hancock, K. & Neill, S. 2005. Reactive oxygen species as signalling molecules. In Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

Epstein, E. 1999. Silicon. *Annual Review of Plant Biology and Plant Molecular Biology*, 50: 641-664.

Epstein, E. & Bloom, A. J. 2005. *Mineral nutrition of plants: Principles and perspectives*. 2nd Ed. Sunderland, MA: Sinauer Associates.

Eraslan, F., Inal, A Savasturk, O. & Gunes, A. 2007. Changes in antioxidative system and membrane damage of lettuce in response to salinity and boron toxicity. *Scientia Horticulturae*, 114:5–10.

Foyer, C.H., Gomez, F.L. & van Heerden, P.D.R. 2005. Glutathione. In Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

Gao, S., Ouyang, C., Wang, S., Xu, T., Tang, L. & Chen, F. 2008. Effects of salt stress on growth, antioxidant enzyme and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedlings. *Plant, Soil Environment*, 54(9): 374–381

Ghassemi F., Jakeman, A.J. & Hix, H.A. 1995. *Salinisation of land and water resources: human causes, extent, management and case studies*. Sydney, Australia: UNSW Press.

Gong, H., Zhu, X., Chen, K., Wang, S. & Zhang, C. 2005. Silicon alleviates oxidative damage of wheat plants in pots under drought. *Plant Science*, 169:313–321.

Harberd, N.P., Belfield, E & Yasumura, Y. 2009. The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: How an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. *Plant Cell*, 21:1328-1339.

Harris, D. 1992. *Hydroponics: The complete guide to gardening without soil*. Cape Town: New Holland Publishing.

Hashemi, A., Abdolzadeh, A. & Sadeghipour, H.R. 2010. Beneficial effects of silicon nutrition in alleviating salinity stress in hydroponically grown canola, *Brassica napus* L., plants. *Soil Science and Plant Nutrition*, 56(2):244-253.

Jaspers, P., Kollist, H., Langebartels, C. & Kangasjärvi, J. 2005.. Plant responses to ozone. In Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

Leyva, R., Sánchez-Rodríguez, E., Ríos, J.J., Rubio-Wilhelmi, M.M., Romero, L., Ruiz, J.M. & Blasco, B. 2011. Beneficial effects of exogenous iodine in lettuce plants subjected to salinity stress. *Plant Science*, 181:195–202.

Liang, T. 1999. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. *Plant and Soil*, 209:217-224.

Liu, X., Ardo, S., Bunning, M., Parry, J., Zhou, K., Stushnoff, C., Stoniker, F. & Yu, L. 2007. Total phenolics content and DPPH radical scavenging activity of lettuce (*Lactuca sativa* L.) grown in Colorado. *LWT – Food Science and Technology*, 40:552-557.

Ma, J.F., Miyake, Y. & Takahashi, E. 2001. Silicon as a beneficial element for crop plants. In Datnoff, L.E., Snyder, G.H. and Korndörfer, G.H. (Eds). *Silicon in agriculture*. New York: Elsevier.

Maas, E.V. & Hoffman, G.J. 1977 . Crop salt tolerance – Current Assesment. *Journal of Irrigation and Drainage Division*,103:115-134.

Magome, H., Yamaguchi, S., Hanada, A., Kamiya, Y. & Oda, K. 2008. The DDF1 transcriptional activator upregulates expression of a gibberellin-deactivating gene, *GA2ox7*, under high-salinity stress in Arabidopsis. *The Plant Journal*, 56(4):613-626.

Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. 2nd Ed. London: Academic Press.

Marshak, S. 2005. *Earth: Portrait of a Planet*. 2nd Ed. New York: W.W. Norton and Company.

Mittler, R (2002). Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*, 7(9):405-410.

Moradi, F. & Ismail, A.M. 2007. Responses of photosynthesis, chlorophyll fluorescence and ROS-scavenging systems to salt stress during seedling and reproductive stages in rice. *Annals of Botany*, 1-13.

Moussa, H.R. 2006. Influence of exogenous application of silicon on physiological response of salt-stressed maize (*Zea mays* L.). *International Journal of Agriculture and Biology*, 8:293-297.

Muckle, M.E. 1993. *Hydroponic nutrients: Easy ways to make your own*. Canada: Growers Press.

Murray, R. K., Granner, D. K., Mayes, P. A., & Rodwell, V. M. 2003. *Harper's Illustrated Biochemistry*. 26th Ed. Columbus, OH: McGraw-Hill.

National Research Council. 1990. *Saline Agriculture: Salt-Tolerant Plants for Developing Countries*. Washington, DC: National Academy Press.

Ndhlala, A.R., Moyo, M. & van Staden, J. 2010. Natural antioxidants: Fascinating or mythical biomolecules? *Molecules*, 15:6905-6930.

Ou, B., Hampsch-Woodill, M. & Prior, R.L. 2001. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *Journal of Agriculture and Food Chemistry*, 49:4619-4626.

Parida, A., Das, A.B. & Das, P. 2002. NaCl stress causes changes in photosynthetic pigments, proteins, and other metabolic components in the leaves of a true mangrove, *Bruguiera parviflora*, in hydroponic cultures. *Journal of Plant Biology*, 24:28-36.

Parida, A.k., Das, A.B., Sanada, Y. & Mohanty, P. 2004. Effects of salinity on biochemical components of the mangrove, *Aegicera corniculatum*. *Aquatic Botany*., 80:77-87.

Resh, H.W. 1995. *Hydroponic food production: A definitive guidebook of soilless food-growing methods*. Santa Barbara, CA: Woodbridge Press.

Romero-Aranda, M.R., Jurado, O. & Cuartero, J. 2006. Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *Journal of Plant Physiology*, 163: 847–855.

Ruiz, J.M. & Blumwald, E. 2002. Salinity-induced glutathione synthesis in *Brassica napus*. *Planta*, 214: 965–969.

- Saqib, M., Zorb, C. & Schubert, S. 2008. Silicon-mediated improvement in the salt resistance of wheat (*Triticum aestivum*) results from increased sodium exclusion and resistance to oxidative stress. *Functional Plant Biology*, 35(7):633-639.
- Savvas, D., Giotis, D., Chatzieustratiou, D., Bakea, M. & Patakioutas, G. 2009. Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environmental and Experimental Botany*, 65(1):11-17.
- Savvas, D., Kizas, G., Lydakis-Simantiris, N., Salahas, G., Papadimitriou, M. & Tsouka N. 2007. Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *European Journal of Horticultural Science*, 72(2):73-79.
- Sekmen, A.H., Türkana, I. & Takiob, S. 2007. Differential responses of antioxidative enzymes and lipid peroxidation to salt stress in salt-tolerant *Plantago maritime* and salt-sensitive *Plantago media*. *Physiologia Plantarum*, 131:399–411.
- Shannon, M.C. & Grieve, C.N. 1999. Tolerance of vegetables to salinity. *Scientia Horticulturae*, 78:5-38.
- Shay, G. 1990. *Saline Agriculture: Salt Tolerant Plants for Developing Countries*. Washington, DC: National Academy Press
- Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.
- Soylemezoglu, G., Demir, K., Inal, A. & Gunes, A. 2009. Effect of silicon on antioxidant and stomatal response of two grapevine (*Vitis vinifera* L.) rootstocks grown in boron toxic, saline and boron toxic-saline soil. *Scientia Horticulturae*, 123(2): 240-246.
- Stern, K.R. 2006. *Introductory plant biology*. 10th Ed. New York: McGraw Hill.
- Tas, G., Papadandonakis, N. & Savvas, D. 2005. Response of lettuce (*Lactuca sativa* L. var. *longifolia*) grown in a closed hydroponic system to NaCl- or CaCl₂-salinity. *Journal of Applied Botany and Food Quality*, 79(2):136-140.
- Tausz, M., Sircelj, H. & Grill, D. 2004. The glutathione system as a stress marker in plant ecophysiology: is a stress response concept valid? *Journal of Experimental Botany*,

55:1855-1862.

Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. & Byrne, D. 2006. Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19:669–675.

Tuna, A.L., Kaya, C., Higgs, D., Murillo-Amador, B., Aydemir, S. & Girgin, A.R. 2008. Silicon improves salinity tolerance in wheat plants. *Environmental and Experimental Botany*, 62(1):10-16.

Urdan, T.C. 2005. *Statistics in Plain English*. 2nd Ed. New Jersey: Lawrence Erlbaum Associates.

Wada, M., Kishikawa, N., Kuroda, N. & Nakashima, K. 2008. Evaluation method for antioxidative activity of health food. In Papadopoulos, K.N. (Ed). *Food Chemistry Research Developments*. New York: Nova Science Publishers.

Wahid, A., Masood, I., Jazed, A. & Rasul, E. 1999. Phenotypic flexibility as marker of sodium chloride tolerance in sunflower genotypes. *Environmental and Experimental Botany*, 42(2):85-94.

Xiong, L. & Zhu, J.K. 2002. Molecular and genetic aspects of plant response to osmotic stress. *Plant, Cell and Environment*, 25:131–139.

Yagi, K. 1984. Assay for plasma lipid peroxides. *Methods in Enzymology*, 109: 328-331.
Yeum, K., Russell, R. and Aldini, G. 2010. Antioxidant activity and oxidative stress: An overview. In Aldini, G., Yeum, K., Niki, E. and Russell, R. (Eds). *Biomarkers for Antioxidant Defense and Oxidative Damage*. Iowa: Blackwell Publishing.

Zhu, Z., Wei, G., Li, J., Qian, Q. & Yu, J. 2004a. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Science*, 167: 527–533.

Zhu, Z., Al-aghabary, K. & Shi, Q. 2004b. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *Journal of Plant Nutrition*, 27(12):2101-2115.




Figure 4.1: Effect of Si applications on Si shoot concentrations in salinity stressed lettuce.




Figure 4.2: Effect of Si applications on Oxidative Radical Absorbance Capacity (ORAC) values in salinity stressed lettuce.




Figure 4.3: Effect of Si applications on Catalase (CAT) concentrations in salinity stressed lettuce.




Figure 4.4: Effect of Si applications on Glutathione (GSH) concentrations in salinity stressed lettuce.

Table 4.1: Effects of Si applications on plant fresh and dry root and shoot weights in salinity stressed lettuce.

NaCl	Si	Fresh Weight (g)		Dry Weight (g)	
		Shoots	Roots	Shoots	Roots
30 mM	0 mM	47.85 ± 17.56	6.49 ± 1.83	2.04 ± 0.51	0.42 ± 0.1
	1 mM	39.1 ± 13.07	5.74 ± 1.81	1.94 ± 0.73	0.34 ± 0.1
	2 mM	43.52 ± 10.49	5.75 ± 1.54	2.12 ± 0.55	0.36 ± 0.09
	4 mM	53.70 ± 11.54	7.64 ± 1.60	2.99 ± 0.56	0.44 ± 0.09
60 mM	0 mM	28.05 ± 6.55	4.52 ± 1.27	1.47 ± 0.31	0.28 ± 0.1
	1 mM	41.73 ± 12.39	7.29 ± 2.38	2.56 ± 0.73**	0.47 ± 0.19*
	2 mM	48.12 ± 14.92*	7.92 ± 2.06**	2.57 ± 0.69**	0.42 ± 0.14
	4 mM	40.97 ± 9.83	6.44 ± 1.52	2.29 ± 0.59	0.40 ± 0.05

Mean values of treated plants were compared to that of the controls (0 mM Si). Significance levels are represented by *P ≤ 0.1, **P ≤ 0.05.

Table 4.2: Effects of Si applications on Total Polyphenolics (TP) and Oxidative Radical Absorbance Capacity, in salinity stressed lettuce.

NaCl	Si	TP (mg GAE g ⁻¹)	ORAC (μmol TE g ⁻¹)
30 mM	0 mM	5.05 ± 0.37	49.28 ± 3.93
	1 mM	4.23 ± 0.35	56.81 ± 4.23
	2 mM	4.71 ± 0.59	54.99 ± 4.07
	4 mM	5.19 ± 0.63	63.16 ± 5.83*
60 mM	0 mM	5.97 ± 0.58	47.72 ± 3.22
	1 mM	5.25 ± 0.59	55.12 ± 9.86
	2 mM	4.98 ± 0.50	54.70 ± 5.74
	4 mM	5.85 ± 0.65	57.51 ± 2.79**

Mean values of treated plants were compared to that of the controls (0 mM Si). Significance levels are represented by *P ≤ 0.1, **P ≤ 0.05, ***P ≤ 0.001.

Table 4.3: Effects of Si applications on Thiobarbituric Acid Reactive Substances (TBARS), Glutathione (GSH), Glutathione disulfide (GSSG), GSH:GSSG ratio and Catalase (CAT), in salinity stressed lettuce.

NaCl	Si	TBARS ($\mu\text{M g}^{-1}$)	GSH ($\mu\text{mol g}^{-1}$)	GSSG ($\mu\text{mol g}^{-1}$)	GSH:GSSG	CAT ($\mu\text{mol min}^{-1} \mu\text{g}^{-1}$)
30 mM	0 mM	0.26 ± 0.03	0.91 ± 0.37	0.15 ± 0.06	5.96 ± 0.81	1.03 ± 0.09
	1 mM	0.26 ± 0.04	0.99 ± 0.2	0.19 ± 0.05	5.20 ± 0.27	0.73 ± 0.09***
	2 mM	0.24 ± 0.03	2.17 ± 0.24***	0.23 ± 0.08	7.94 ± 0.59*	0.86 ± 0.12
	4 mM	0.29 ± 0.04	1.43 ± 0.46	0.24 ± 0.07	5.91 ± 0.67	0.77 ± 0.11**
60 mM	0 mM	0.25 ± 0.02	1.04 ± 0.83	0.15 ± 0.1	6.94 ± 1.76	1.12 ± 0.10
	1 mM	0.28 ± 0.03	2.14 ± 0.37**	0.29 ± 0.07*	7.69 ± 1.87	1.03 ± 0.06
	2 mM	0.25 ± 0.03	1.47 ± 0.38	0.25 ± 0.06	5.87 ± 0.15	0.95 ± 0.11
	4 mM	0.26 ± 0.03	2.07 ± 0.28**	0.30 ± 0.05**	7.06 ± 1.01	1.28 ± 0.16

Mean values of treated plants were compared to that of the controls (0 mM Si). Significance levels are represented by *P ≤ 0.1, **P ≤ 0.05, ***P ≤ 0.001.

CHAPTER FIVE

GENERAL DISCUSSION AND CONCLUSION

Silicon, in this study, and in the myriad of other studies investigating silicon's ability to ameliorate abiotic and biotic stress factors, has proved its status as a quasi-essential nutrient in plants. With salinisation being a widespread occurrence in many soils, and the accumulation of salts in soilless growing systems a regular incident, Si shows a remarkably cost efficient and environmentally friendly way of mitigating the deleterious effects that are associated with salinity stress in plants, ultimately increasing yield.

The studies herein confirm the beneficial effects of Si with regards to NaCl salinity stress, with the first study reporting on yield, chlorophyll, leaf numbers, height and elemental content of lettuce plants. Although leaf number, plant height and chlorophyll content showed limited or no significance when comparing the treated plants to the controls, fresh root and shoot weights showed clear increases when compared to the controls grown without Si, as did dry root and shoot weights.

Most noticeable, and in agreement with the majority of other studies evaluating Si and NaCl stress, Na tissue content showed a clear tendency to decrease with applications of Si, and this can be seen as the foremost reason for the increase in yield and therefore the mitigating properties associated with Si applications; the Si-Na interaction. Although the mechanism of this Si-Na interaction has only been the subject of a limited number of studies, it is probable that it is the result of Na dilution due to an increase in water use efficiency and therefore a decrease in Na toxicity (Romero-aranda et al., 2005), although data from this study cannot support this statement. This is supported by studies reporting positive effects of Si applications on drought stressed plants.

The antioxidant status and oxidative damage study showed positive results with applications of Si, although the interpretation of these results were subject to a higher degree of speculation than the more obvious growth parameters of the first study. Concentrations of both GSH and GSSG tended to increase with applications of Si. TBARS concentrations showed a surprisingly limited trend to increase or decrease with or without applications of Si, a decrease of which would have given a clear indication of oxidative damage mitigation. Most noticeable with regards to this study was the tendency for CAT activities to decrease with applications of Si. CAT is a major ROS scavenging enzyme seen as indispensable, with stress bringing about the production of CAT (Mittler, 2002). It follows that with increased CAT production during stress, a decrease of CAT would be a sign of less oxidative stress, as was

found with applications of Si in this study. Contrarily, and as noticeable, ORAC values increased with Si applications, indicating a higher capacity to scavenge ROS, and therefore a beneficial characteristic brought about by Si.

From these results, it can be concluded that Si has a significantly positive effect on yield, antioxidant status, oxidative damage and NaCl stress mitigation as a whole. However, the connection between the resultant antioxidant status to yield data, showed no correlation. Additionally, oxidative damage was somewhat unconvincing, therefore additions of higher concentrations of NaCl would possibly lead to more conclusive oxidative damage data, and perhaps other non-antioxidant related factors as well. These factors are important to consider for future studies.

REFERENCES

- Aebi, H. 1984. Catalase in vitro. *Methods in Enzymology*, 105:121–126.
- Ahmad, R., Zaheer, S.H. and Ismail, S. 1992. Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Science*, 85(1):43-50.
- Arnon, D.L. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidases in *Beta vulgaris*. *Plant Physiology*. 24:1-15.
- Arnon, D.L. & Stout, P.R. 1939. The essentiality of certain elements in minute quantities with special reference to copper. *Plant Physiology* 14: 371-375.
- Asensi, M., Sastre, J., Pallardo, F.V., Lloret, A., Lehner, M., Garcia-de-la Asuncion, J. & Viña J. 1999. Ratio of reduced to oxidised glutathione as indicator of oxidative stress status and DNA damage. *Methods in Enzymology*. 1999(299):267-76.
- Ashraf, M., Rahmatullah., Afzal, M., Ahmed, R., Mujeeb, F., Sarwar, A. & Ali, L. 2010. Alleviation of detrimental effects of NaCl by silicon nutrition in salt-sensitive and salt-tolerant genotypes of sugarcane (*Saccharum officinarum* L.). *Plant and Soil*, 326(1):381-391.
- Barroso, M.C. & Alvarez, C.E. 1997. Toxicity symptoms and tolerance of strawberry to salinity in the irrigation water. *Scientia Horticulturae.*, 71:177-188.
- Belanger, R.R, Bowen, P.A., Ehret, D.L. & Menzies, J.G. 1995. Soluble silicon: it's role in crop and disease management of greenhouse crops. *Plant Disease*, 79(4):329-336.
- Bor, M., Özdemir, F. & Türkan, I. 2003. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Science*, 164:77-84.
- Bradbury, M. & Ahmad, R. 1990. The effect of silicon on the growth of *Prosopis juliflora* growing in saline soil. *Plant and Soil*, 125:71-74.
- Carter, D.L. 1981. Salinity and plant productivity. *CRC Handbook of Agricultural Productivity*, 1:117-133.

- Dat, J., Vandenberghe, S., Vranová, E., Van Montagu, M., Inzé, D. & Van Breusegem, F. 2000. Dual action of the active oxygen species during plant stress responses. *Cellular and Molecular Life Sciences*, 57:779–795.
- Desikan, R., Hancock, K. & Neill, S. 2005. Reactive oxygen species as signalling molecules. In Smirnov, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.
- Epstein, E. 1994. The anomaly of silicon in plant biology. *Proceedings of the National Academy of Sciences of the United States of America*, 91:11-17.
- Epstein, E. 1999. Silicon. *Annual Review of Plant Biology and Plant Molecular Biology*, 50: 641-664.
- Epstein, E. & Bloom, A. J. 2005. *Mineral nutrition of plants: Principles and perspectives*. 2nd Ed. Sunderland, MA: Sinauer Associates.
- Eraslan, F., Inal, A Savasturk, O. & Gunes, A. 2007. Changes in antioxidative system and membrane damage of lettuce in response to salinity and boron toxicity. *Scientia Horticulturae*, 114:5–10.
- Feierabend, J. 2005. Catalase in plants: molecular and functional properties and role in stress defence. In Smirnov, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.
- Flowers, T.J. & Yeo, A.R. 1981. Variability in the resistance of sodium chloride salinity within rice (*Oryza sativa* L.) varieties. *New Phytology*, 88:363-373.
- Foyer, C.H., Gomez, F.L. & van Heerden, P.D.R. 2005. Glutathione. In Smirnov, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.
- Gao, S., Ouyang, C., Wang, S., Xu, T., Tang, L. & Chen, F. 2008. Effects of salt stress on growth, antioxidant enzyme and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedlings. *Plant, Soil Environment*, 54(9): 374–381
- Ghassemi F., Jakeman, A.J. & Hix, H.A. 1995. *Salinisation of land and water resources: human causes, extent, management and case studies*. Sydney, Australia: UNSW Press.

Gong, H., Zhu, X., Chen, K., Wang, S. & Zhang, C. 2005. Silicon alleviates oxidative damage of wheat plants in pots under drought. *Plant Science*, 169:313–321.

Grace, S.C. 2005. Phenolics as antioxidants. In Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

Hadfield, J. 2001. *The A-Z of vegetable gardening in South Africa*. Cape Town: Struik.

Halliwell, B & Gutteridge, J.M.C. 1999. *Free Radicals in Biology and Medicine*. 3rd Ed. London: Oxford UP.

Han, H.S. & Lee, K.D. 2005. Plant growth promoting rhizobacteria effect on antioxidant status, photosynthesis, mineral uptake and growth of lettuce under soil salinity. *Research Journal of Agriculture and Biological Sciences*, 1:210-215.

Harberd, N.P., Belfield, E & Yasumura, Y. 2009. The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: How an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. *Plant Cell*, 21:1328-1339.

Hashemi, A., Abdolzadeh, A. & Sadeghipour, H.R. 2010. Beneficial effects of silicon nutrition in alleviating salinity stress in hydroponically grown canola, *Brassica napus* L., plants. *Soil Science and Plant Nutrition*, 56(2):244-253.

Harris, D. 1992. *Hydroponics: The complete guide to gardening without soil*. Cape Town: New Holland Publishing.

Hemy, C. 1984. *Growing vegetables in South Africa*. Johannesburg: Macmillan South Africa.

Hiscox, J.D. & Israelstam, G.F. 1979. A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany*, 57:1332-1334.

Hitchem, H., Mounir, D. & Naceur E.A. 2009. Differential responses of two maize varieties to salt stress: Changes in polyphenols composition of foliage and oxidative damages. *Industrial Crops and Products*, 30:144-151.

Jaspers, P., Kollist, H., Langebartels, C. & Kangasjärvi, J. 2005.. Plant responses to ozone. In Smirnoff, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

- Koolman, J & Rhoem, K. 2005. Color Atlas of Biochemistry. 2nd Ed. Stuttgart, Germany: Thieme.
- Lerner R.E., Meacham, S. & Burns. 1993. *Western Civilizations*. United States of America: W.W. Norton & Company.
- Leyva, R., Sánchez-Rodríguez, E., Ríos, J.J., Rubio-Wilhelmi, M.M., Romero, L., Ruiz, J.M. & Blasco, B. 2011. Beneficial effects of exogenous iodine in lettuce plants subjected to salinity stress. *Plant Science*, 181:195–202.
- Liang, T. 1999. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. *Plant and Soil*, 209:217-224.
- Liang, Y., Chen, Q., Liu, Q., Zhang, W. & Ding, R. 2003. Exogenous silicon increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley. *Journal of Plant Physiology*, 160(10):1157-1164.
- Liu, X., Ardo, S., Bunning, M., Parry, J., Zhou, K., Stushnoff, C., Stoniker, F. & Yu, L. 2007. Total phenolics content and DPPH radical scavenging activity of lettuce (*Lactuca sativa* L.) grown in Colorado. *LWT – Food Science and Technology*, 40:552-557.
- Ma, J.F., Miyake, Y. & Takahashi, E. 2001. Silicon as a beneficial element for crop plants. In: Datnoff L.E., Snyder, G.H. and Korndorf G.H. *Silicon in Agriculture*. Amsterdam: Elsevier Science.
- Maas, E.V. & Hoffman, G.J. 1977 . Crop salt tolerance – Current Assesment. *Journal of Irrigation and Drainage Division*,103:115-134.
- Magome, H., Yamaguchi, S., Hanada, A., Kamiya, Y. & Oda, K. 2008. The DDF1 transcriptional activator upregulates expression of a gibberellin-deactivating gene, *GA2ox7*, under high-salinity stress in *Arabidopsis*. *The Plant Journal*, 56(4):613-626.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. 2nd Ed. London: Academic Press.
- Marshak, S. 2005. *Earth: Portrait of a Planet*. 2nd Ed. New York: W.W. Norton and Company.
- Mittler, R (2002). Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*, 7(9):405-410.

Moradi, F. & Ismail, A.M. 2007. Responses of photosynthesis, chlorophyll fluorescence and ROS-scavenging systems to salt stress during seedling and reproductive stages in rice. *Annals of Botany*, 1-13.

Moussa, H.R. 2006. Influence of exogenous application of silicon on physiological response of salt-stressed maize (*Zea mays* L.). *International Journal of Agriculture and Biology*, 8:293-297.

Muckle, M.E. 1993. *Hydroponic nutrients: Easy ways to make your own*. Canada: Growers Press.

Murray, R. K., Granner, D. K., Mayes, P. A., & Rodwell, V. M. 2003. *Harper's Illustrated Biochemistry*. 26th Ed. Columbus, OH: McGraw-Hill.

National Research Council. 1990. *Saline Agriculture: Salt-Tolerant Plants for Developing Countries*. Washington, DC: National Academy Press.

Ndhlala, A.R., Moyo, M. & van Staden, J. 2010. Natural Antioxidants: Fascinating or Mythical Biomolecules? *Molecules*, 15: 6905-6930.

Ou, B., Hampsch-Woodill, M. & Prior, R.L. 2001. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *Journal of Agriculture and Food Chemistry*, 49:4619-4626.

Parida, A., Das, A.B. & Das, P. 2002. NaCl stress causes changes in photosynthetic pigments, proteins, and other metabolic components in the leaves of a true mangrove, *Bruguiera parviflora*, in hydroponic cultures. *Journal of Plant Biology*, 24:28-36.

Parida, A.k., Das, A.B., Sanada, Y. & Mohanty, P. 2004. Effects of salinity on biochemical components of the mangrove, *Aegicera corniculatum*. *Aquatic Botany.*, 80:77-87.

Reezi, S., Babalar, M. & Kalantari, S. 2009. Silicon alleviates salt stress, decreases malondialdehyde content and affects petal color of salt stressed cut rose (*Rosa xhybrida* L.) 'Hot Lady'. *African Journal of Biotechnology*, 8(8):847-855.

Resh, H.W. 1995. *Hydroponic food production: A definitive guidebook of soilless food-growing methods*. Santa Barbara, CA: Woodbridge Press.

- Romero-Aranda, M.R., Jurado, O. & Cuartero, J. 2006. Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *Journal of Plant Physiology*, 163: 847–855.
- Ruiz, J.M. & Blumwald, E. 2002. Salinity-induced glutathione synthesis in *Brassica napus*. *Planta*, 214: 965–969.
- Rush, D.W. & Epstein, E. 1976. Differences between salt-sensitive and salt-tolerant genotypes of the tomato. *Plant Physiology*, 57: 162-166.
- Saqib, M., Zorb, C. & Schubert, S. 2008. Silicon-mediated improvement in the salt resistance of wheat (*Triticum aestivum*) results from increased sodium exclusion and resistance to oxidative stress. *Functional Plant Biology*, 35(7):633-639.
- Savvas, D., Giotis, D., Chatzieustratiou, D., Bakea, M. & Patakioutas, G. 2009. Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environmental and Experimental Botany*, 65(1):11-17.
- Savvas, D., Kizas, G., Lydakis-Simantiris, N., Salahas, G., Papadimitriou, M. & Tsouka N. 2007. Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *European Journal of Horticultural Science*, 72(2):73-79.
- Sekmen, A.H., Türkana, I. & Takiob, S. 2007. Differential responses of antioxidative enzymes and lipid peroxidation to salt stress in salt-tolerant *Plantago maritime* and salt-sensitive *Plantago media*. *Physiologia Plantarum*, 131:399–411.
- Shannon, M.C. & Grieve, C.N. 1999. Tolerance of vegetables to salinity. *Scientia Horticulturae*, 78:5-38.
- Shay, G. 1990. *Saline Agriculture: Salt Tolerant Plants for Developing Countries*. Washington, DC: National Academy Press
- Soylemezoglu, G., Demir, K., Inal, A. & Gunes, A. 2009. Effect of silicon on antioxidant and stomatal response of two grapevine (*Vitis vinifera* L.) rootstocks grown in boron toxic, saline and boron toxic-saline soil. *Scientia Horticulturae*, 123(2): 240-246.
- Smirnov, N. (Ed). *Antioxidants and Reactive Oxygen Species in Plants*. Chennai, India: Blackwell Publishing.

- Stern, K.R. 2006. *Introductory plant biology*. 10th Ed. New York: McGraw Hill.
- Tas, G., Papadandonakis, N. & Savvas, D. 2005. Response of lettuce (*Lactuca sativa* L. var. *longifolia*) grown in a closed hydroponic system to NaCl- or CaCl₂-salinity. *Journal of Applied Botany and Food Quality*, 79(2):136-140.
- Tausz, M., Sircelj, H. & Grill, D. 2004. The glutathione system as a stress marker in plant ecophysiology: is a stress response concept valid? *Journal of Experimental Botany*, 55:1855-1862.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L. & Byrne, D. 2006. Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19:669–675.
- Tuna, A.L., Kaya, C., Higgs, D., Murillo-Amador, B., Aydemir, S. & Girgin, A.R. 2008. Silicon improves salinity tolerance in wheat plants. *Environmental and Experimental Botany*, 62(1):10-16.
- Urdan, T.C. 2005. *Statistics in Plain English*. 2nd Ed. New Jersey: Lawrence Erlbaum Associates.
- Wada. M., Kishikawa, N., Kuroda, N. & Nakashima, K. 2008. Evaluation method for antioxidative activity of health food. In Papadopoulos, K.N. (Ed). *Food Chemistry Research Developments*. New York: Nova Science Publishers.
- Wang, X., Wei, Z., Liu, D. & Zhao, G. 2011. Effects of NaCl and silicon on activities of antioxidative enzymes in roots, shoots and leaves of alfalfa. *African Journal of Biotechnology*, 10:545-549.
- Xiong, L. & Zhu, J.K. 2002. Molecular and genetic aspects of plant response to osmotic stress. *Plant, Cell and Environment*, 25:131–139.
- Yagi, K. 1984. Assay for plasma lipid peroxides. *Methods in Enzymology*, 109: 328-331.
- Yan-fang, R. and Jun-yu, H. 2008. Effects of NaCl stress on growth and photosynthesis characteristics of lettuce (*Lactuca sativa* L.) seedlings. *Acta Agriculturae Boreali-Sinica*, 4.

Yeum, K., Russell, R. and Aldini, G. 2010. Antioxidant activity and oxidative stress: An overview. In Aldini, G., Yeum, K., Niki, E. and Russell, R. (Eds). *Biomarkers for Antioxidant Defense and Oxidative Damage*. Iowa: Blackwell Publishing.

Wahid, A., Masood, I., Jazed, A. & Rasul, E. 1999. Phenotypic flexibility as marker of sodium chloride tolerance in sunflower genotypes. *Environmental and Experimental Botany*, 42(2):85-94.

Zhu, Z., Wei, G., Li, J., Qian, Q. & Yu, J. 2004a. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Science*, 167: 527–533.

Zhu, Z., Al-aghabary, K. & Shi, Q. 2004b. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *Journal of Plant Nutrition*, 27(12):2101-2115.

APPENDIX A: The subirrigative, closed soilless growing systems for the cultivation of *Lactuca sativa* L. 'Eish!'.



Lettuce seedlings 1 week after transplanting in the soilless growing systems.



Figure 3.7. Lettuce seedlings 4 week after transplanting in the soilless growing systems.

Full Length Research Paper

The alleviation of salinity induced stress with applications of silicon in soilless grown *Lactuca sativa* L. 'Eish!'

Milne C. J.¹, Laubscher C. P.^{1*} and Ndakidemi P. A.²

¹Faculty of Applied Science, Cape Peninsula University of Technology, Cape Town Campus, Keizersgracht, P. O. Box 652, Cape Town 8000, South Africa.

²The Nelson Mandela African Institute of Science and Technology, P. O. Box 447, Arusha-Tanzania.

Accepted 5 December, 2011

The effects of 30 and 60 mM NaCl on Lettuce (*Lactuca sativa* L. 'Eish!'), grown in soilless culture, with additions of 0, 1, 2 and 4 mM Si was evaluated. Height, leaf number, weight, chlorophyll content and elemental analysis of plants were examined. Fresh root and shoot weight significantly increased with additions of 2 mM Si compared to that of the 60 mM NaCl control, with notable increases with Si treatments of 1 and 4 mM. Dry root and shoot weight with additions of 1 mM Si, and in shoot weight with additions of 2 mM Si, significantly increased when compared to the 60 mM NaCl control. Increases in dry root and shoot weight of the other Si additions lead to clear, yet insignificant increases in dry weights. Few differences in Cl content were evident amongst treatments, besides a significant increase in root content and a significant decrease in shoot content with applications of 4 mM Si exposed to 60 mM NaCl. Na shoot content of 1, 2 and 4 mM Si at 60 mM NaCl showed a significant decrease compared to that of the control. Applications of 2 and 4 mM Si showed increased shoot Si when compared to both the 30 and 60 mM NaCl controls. With clear evidence for a reduction in Na content in shoots with applications of Si at higher NaCl concentrations, and further increases in plant weights, it can be stated that applications of Si are of a distinct benefit for the growth of lettuce.

Key words: Lettuce, Si, NaCl, salt, sodium chloride, hydroponics.

INTRODUCTION

Silicon [Si], an element whose abundance in the earth's crust is second only to oxygen (Marshak, 2005), can contribute as much as 0.1 to 10% of the dry matter of plants. These levels are equal to or exceeding those of essential macro nutrients (Epstein and Bloom, 2005). Si has in the past been classified as a non-essential element for plant nutrition, that is, one which does not meet the criteria for essentiality in the classic definition of essentiality proposed by Arnon and Stout (1939). More recently Si has been classified as either a beneficial element (Marschner, 1995), or as a quasi-essential

element in plants (Epstein, 1999).

Contributing to its classification as a beneficial or quasi-essential element in plants is its role as a plant stress ameliorator (Epstein, 1994, 1999; Belanger et al., 1995). An important aspect of this is the ability of Si to ameliorate salt toxicity, defined as conditions of high salt build-up, with sodium chloride [NaCl] being the most common cause of saline conditions (Harris, 1992; Epstein and Bloom, 2005).

Salinity has two methods of causing abnormalities in plant growth. The first, which is independent of the type of salt, is by creating a higher osmotic potential in the medium surrounding the roots, and because osmosis is the principle method of water absorption in plants (Stern, 2006), it results in a decrease in plant-water availability (Carter, 1981). The second, which is salt dependent, is

*Corresponding author. E-mail: laubscherc@cput.ac.za. Tel: +27 21 4603198. Fax: +27 21 460 3193.

caused by an excess of available ions. In the case of NaCl salinity, the ions in question are sodium [Na⁺] and chloride [Cl⁻]. Cl is known to be an essential micronutrient in plants (Epstein and Bloom, 2005; Stern, 2006), and some argued that Cl should be classified as a macro-nutrient, contributing as much as 1% of the dry weight of some plants (Muckle, 1993). Na, on the other hand, is required by only a few plants (Epstein and Bloom, 2005), and has been shown to be the primary cause of growth suppression in lettuce (Tas et al., 2005). In general, high concentrations of both Na and Cl are associated with leaf scorching (Shannon and Grieve, 1999), necrosis and defoliation (Maas and Hoffman, 1977).

The mitigation of NaCl induced toxicity with the application of Si has been demonstrated in many important agricultural and ornamental crops, including Alfalfa (Wang et al., 2011), tomato (Romero-Aranda et al., 2006), wheat (Ahmad et al., 1992), zucchini (Savvas et al., 2009), grapevine (Soylemezoglu et al., 2009), mesquite (Bradbury and Ahmad, 1990), roses (Savvas et al., 2007; Reezi et al., 2009) and several others. The present study investigated the ability of Si to alleviate NaCl toxicity in lettuce (*Lactuca sativa* L. 'Eish!').

MATERIALS AND METHODS

The experiment took place within the experimental greenhouse located at the Cape Peninsula University of Technology in Cape Town, South Africa - 33° 55' 58.27''S, 18° 25' 57.04' E. The temperature within the greenhouse ranged from 9 to 29°C, the mean being 17.6°C and the humidity ranged from 44 to 99%, the mean being 77%.

L. sativa L. 'Eish!' seeds (Hygrotech, South Africa) were sown in vermiculite, and germinated under 40% shade-cloth. Once all seeds had germinated (8 days after sowing) they were fertilised daily with half strength CHEMICULT (Starke Ayres, South Africa) nutrient solution, and increased gradually to full strength (Harris, 1992). Once the seedlings were fully established (40 days after sowing) 80 plants were randomly selected and transplanted into individual 12.5 cm pots, containing expanded clay as the medium, and placed within the soilless growing system. The sub-irrigation, closed, soilless growing system consisted of a nutrient solution holding tray, delivering a constant ± 1 cm of circulating nutrient solution to each treatment. Using a randomised block design, 10 plants were placed within eight individual soilless culture units, allowing for 10 repeats of each.

Basal nutrient solution

Analytical grade, potassium nitrate [KNO₃], calcium nitrate [(Ca(NO₃)₂·4H₂O)], ammonium dihydrogen phosphate [NH₄H₂PO₄], and magnesium sulphate [MgSO₄·7H₂O], were used to add 16 mM N, 6 mM K, 4 mM Ca, 2 mM P, 1 mM S and 1 mM Mg to the nutrient solution, which was based on the modified Hoagland solution presented by Epstein and Bloom (2005). Micro nutrients were supplied by using HYGROPLEX (Hygrotech, South Africa), adding 58 µM Boron [B], 11.4 µM manganese [Mn], 7.6 µM zinc [Zn], 1 µM copper [Cu], 0.65 µM molybdenum [Mo] and 31.9 µM iron [Fe] to the nutrient solution. The complete basal nutrient solution was prepared using deionised water and had an electrical conductivity (EC) of 1.9 mS cm⁻¹.

Silicon and NaCl treatments

Four levels of Si, 0, 1, 2 and 4 mM, were supplied by adding AGRISIL K50 (PQ Silicas, South Africa) - Potassium silicate [K₂SiO₃], each of which were combined with two levels of NaCl, 30 and 60 mM, making a total of 8 treatments. Treatments, which were applied in a single dose, commenced one week after transplantation. Newly prepared nutrient solutions containing 30 mM NaCl had a mean EC of 5.20 mS cm⁻¹ and solutions containing 60 mM NaCl had a mean EC of 8.10 mS cm⁻¹. These EC levels were maintained daily. Potassium [K] was balanced in all treatments by subtracting additional K from KNO₃, and supplementing lost Nitrate [NO₃] by adding nitric acid [HNO₃]. The nutrient solution, which was replaced weekly, was maintained daily at a pH of 6.0 using hydrochloric acid [HCl] and sodium hydroxide [NaOH].

Plant growth parameters

Plant growth, in terms of plant height and leaf number, of all treatments, were measured at weekly intervals. Plant height was measured with a measuring tape, from the surface of the medium to the tip of the tallest leaf.

Five plants per treatment were harvested after 11 weeks from sowing (Hadfield, 2001), and separated into roots and shoots. Roots were rinsed once with tap water and twice with deionised water before being patted dry with paper towel. After the fresh weight had been recorded, plants were immediately placed within a specimen oven at 60°C until constant weight, and the dry weight recorded.

Determination of chlorophyll content in plant leaves

Extraction of chlorophyll by dimethyl sulphoxide [DMSO] was done following a modified method described by Hiscox and Israelstam (1979). This method allows for the extraction of chlorophyll without plant tissue maceration. Five lettuce plants per treatment were lyophilised and ground to a fine powder using a mortar and pestle and stored at -80°C until analysed. 100 mg of the powder was placed in a 15 ml vial containing 7 ml DMSO and incubated at 65°C for 20 min. After the incubation period, an additional 3 ml of DMSO was added to the vial, and stored at 4°C. After a 24 h storage period, the extract was centrifuged at 3220 g for 10 min, after which the supernatant was transferred to a new 15 ml vial. Before spectrophotometrical analysis, the supernatant was diluted to a factor of 15 of which 1 ml was placed in a cuvette and absorbance values of 645 and 663 nm recorded. Absorbance values were used in the equation proposed by Arnon (1949), to determine total leaf chlorophyll content against DMSO blank, expressed as mg g⁻¹ dry weight.

Elemental analysis

Dried plant material was used to analyse for Na, Cl and Si. Na and Si was analysed by ashing the ground sample and dissolving the ash in HCl. Elemental concentrations were determined using an inductively-coupled plasma (ICP) emission spectrophotometer. Cl was analysed by digesting the dried, ground sample with HNO₃. An excessive quantity of silver nitrate [AgNO₃] was added, followed by back titration with potassium thiocyanate [KSCN]. Elemental concentrations of Na, Cl and Si are expressed as mg g⁻¹ of dried plant material (A. Van Deventer, Bemlab, South Africa, Personal Communication).

Statistical analysis

Statistical analysis was performed by using two-way independent




Figure 1. The effect of 30 and 60 mM NaCl on plant height, with additions of 0, 1, 2 and 4 mM Si. Measurements were taken weekly from 27 May (1) to 17 June (4) 2011.




Figure 2. The effect of 30 and 60 mM NaCl on fresh root weight, with additions of 0, 1, 2 and 4 mM Si.

analysis of variance (ANOVA) followed by the Bonferroni post test, with Si and NaCl concentrations being the factors assessed. Computations were executed with the software program SPSS (Urda, 2005).

RESULTS

Leaf number showed no significant differences between Si treatments, whether compared to the control, containing 0 mM Si, or to each other (data not shown). Plant height showed similar insignificance between treatments and controls, besides the expected general increase in height over the growing period (Figure 1). There was a

significant interaction between salt concentration and silicon treatments on fresh root weight ($P \leq 0.05$) and fresh shoot weight ($P \leq 0.1$). Main effect analysis showed a significant decrease in fresh root weight and fresh shoot weight ($P \leq 0.1$ and $P \leq 0.05$ respectively) between the two controls (Figures 2 and 3). Fresh root weight and fresh shoot weight showed no significant differences at Si levels of 1 and 4 mM whether grown with 30 or 60 mM NaCl, when compared to their respective control. The treatment of 2 mM Si exposed to 60 mM did however show a significant increase in fresh root ($P \leq 0.05$) and shoot weight ($P \leq 0.1$) when compared to the control. Although only 2 mM Si showed a significant effect on




Figure 3. The effect of 30 and 60 mM NaCl on fresh shoot weight, with additions of 0, 1, 2 and 4 mM Si.




Figure 4. The effect of 30 and 60 mM NaCl on dry root weight, with additions of 0, 1, 2 and 4 mM Si.

fresh root and shoot weight, there was a marked, yet insignificant, increase in fresh root and shoot weight of 4 mM Si at 30 mM NaCl, and 1 mM and 4 mM Si at 60 mM when compared to the control (Figures 2 and 3).

There was also a significant interaction between salt concentration and silicon treatments on dry root weight ($P \leq 0.1$) and dry shoot weight ($P \leq 0.05$). Main effect analysis showed that dry root weight significantly decreased ($P \leq 0.1$) between the control of 30 mM NaCl and that of 60 mM NaCl. Besides a significant increase ($P \leq 0.1$) in dry root weight between 1 mM Si exposed to 60 mM NaCl and that of the control, the other treatments

showed no significant differences, even though there was again a marked increase in weight, especially when compared to the 60 mM NaCl control (Figure 4). Dry shoot weight showed no significant increase between controls. However, there were significant increases ($P \leq 0.05$) in weight of 1 and 2 mM Si when compared to the 60 mM NaCl control. Again, although other treatments resulted in insignificant differences, there was a marked increase in root and shoot dry weight when compared to that of the control (Figure 5).

Although there was a significant interaction between salt concentration and silicon treatments on root ($P \leq 0.001$)




Figure 5. The effect of 30 and 60 mM NaCl on dry shoot weight, with additions of 0, 1, 2 and 4 mM Si.

Table 1. The effect of 30 and 60mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4mM Si.

Treatment		Cl (mg g^{-1})		Na (mg g^{-1})		Si (mg g^{-1})	
NaCl (mM)	Si (mM)	Root	Shoot	Root	Shoot	Root	Shoot
30	0	12.18	16.78	7.17	11.11	0.42	0.06
	1	9.78	18.70	8.12	11.41	0.44	0.12
	2	9.60	22.54	8.59	11.22	0.51	0.15**
	4	14.42	13.74	8.10	11.64	0.43	0.43***
60	0	15.98	21.20	14.30	16.88	0.37	0.07
	1	14.82	13.62	9.43	13.75***	0.29	0.11
	2	7.66	14.20	10.77	15.40**	0.45	0.16***
	4	74.8***	12.74*	8.83	14.00***	0.40	0.18***

and shoot ($P \leq 0.05$) Cl content, main effect analysis showed that Cl concentrations only increased significantly in roots of the 4 mM Si with 60 mM NaCl treated plants and the shoots of 2 mM Si plants. This in turn corresponded to a significant decrease ($P \leq 0.1$) in shoot Cl when compared to the control at 60 mM NaCl (Table 1). When comparing Cl content of plants treated with the same level of Si, 2 mM Si at 60 mM NaCl showed a significant decrease in shoot Cl content ($P \leq 0.05$) compared to the plants treated with 30 mM NaCl. Contrarily, plants treated with 4 at 60 mM NaCl showed a significant increase in root Cl content ($P \leq 0.01$) when compared to the plants treated with 30 mM NaCl (Table 2).

There was a significant interaction between salt concentration and silicon treatments on shoot Na content ($P \leq 0.001$) but not on root Na content. Although no significant differences in root Na content were obtained, there was a marked decrease in Na root concentration at 60 mM NaCl (Figure 6). Main effect analysis showed that Na shoot content at 30 mM NaCl mirrored the results of


that of the roots, and showed no significant differences between treatments. However, Na shoot content at 60 mM NaCl with Si levels of 1, 2 and 4m M all resulted in a significant decrease in shoot Na content compared to that of the control (Table 1). When comparing Na content of plants treated with the same level of Si, root Na significantly increase between the controls of 30 and 60 mM NaCl ($P \leq 0.001$) and shoot Na significantly increase ($P \leq 0.001$) with 60 mM NaCl when compared to 30 mM NaCl at all Si levels (Table 2).

There was a significant interaction between salt concentration and silicon treatments on Si shoot content ($P \leq 0.001$), but not on Si root content. Main effect analysis showed that there was however a significant decrease ($P \leq 0.05$) of root Si between treatments at 30 and 60 mM NaCl receiving 1 mM Si. Si shoot concentrations of 2 and 4 mM, at both 30 and 60 mM NaCl showed a significant increase when compared to the control. When the two NaCl levels at the same 4 mM Si treatment were compared, a significant decrease ($P \leq 0.01$)

Table 2. The effect of 30 and 60 mM NaCl on Cl, Na and Si root and shoot content, with additions of 0, 1, 2 and 4 mM Si.

Treatment		Cl (mg g ⁻¹)		Na (mg g ⁻¹)		Si (mg g ⁻¹)	
Si (mM)	NaCl (mM)	Roots	Shoots	Roots	Shoots	Roots	Shoots
0	30	12.18	16.78	7.17	11.11	0.42	0.06
	60	15.98	21.20	14.30***	16.88***	0.37	0.07
1	30	9.78	18.70	8.12	11.41	0.44	0.12
	60	14.82	13.62	9.43	13.75***	0.29**	0.11
2	30	9.60	22.54	8.59	11.22	0.51	0.15
	60	7.66	14.20**	10.77	15.40***	0.45	0.16
4	30	14.42	13.74	8.10	11.64	0.43	0.43
	60	74.8***	12.74	8.83	14.00***	0.40	0.18***

Values presented are means \pm SE, n = 5. *, **, *** = significant at $P \leq 0.1^*$, $P \leq 0.05^{**}$ or $P \leq 0.01^{***}$. Significance compared to that of the same Si concentrations, with different NaCl concentrations.

**Figure 6.** The effect of 30 and 60 mM NaCl on Na root content, with additions of 0, 1, 2 and 4 mM Si.

in Si was observed in the shoots. This was apparent in the 4 mM treatment alone (Table 2).

There was a significant interaction between salt concentration and silicon treatments on chlorophyll content ($P \leq 0.05$). Main effect analysis revealed that, when comparing Si treated plants to the controls, there was only a significant decrease in chlorophyll with applications of 2 mM Si at 30 mM to that of the control (Figure 7).

DISCUSSION

The results obtained from this study support the

fundamental findings of previous studies, that Si has beneficial mitigating properties with regards to NaCl induced salinity stress. These findings indicate some benefits of Si applications at 30 mM NaCl; however, the benefits of Si seem to be more apparent with the higher experimental concentrations of 60 mM NaCl. This is evident when looking at the significant increase in fresh root and shoot weight at 2 mM Si at 60 mM NaCl, the significant increases in dry root, at 1 mM Si, dry shoot weight at 1 and 2 mM, and the overall marked increases in fresh and dry weights when comparing all Si treatments to that of the 60 mM NaCl control (Figures 2 to 5). These findings are supported by the literature which




Figure 7. The effect of 30 and 60 mM NaCl on total chlorophyll content with additions of 0, 1, 2 and 4 mM Si.

states that dry shoot and root weights of plants increase with applications of Si compared to a range of NaCl concentrations lacking Si (Romero-Aranda et al., 2006; Tuna et al., 2008; Ashraf et al., 2010).

Chlorophyll content showed no significant differences, with only one exception with an application 2 mM Si at 30 mM NaCl (Figure 7). This is in line with Tas et al. (2005), who although did not investigate Si, reported negligible effects of NaCl on chlorophyll content on lettuce. However, in a study conducted by Hashemi et al. (2010), evaluating the mitigating effects of Si on Canola, a vast increase in chlorophyll content was found.

A notable oddity with regards to elemental analysis was the exceptional accumulation of Cl in the roots of the plants treated with 4 mM Si at 60 mM (Tables 1 and 2). This, when compared to the control, corresponded to a significant decrease in Cl accumulation in shoots. Savvas et al. (2007, 2009) found similar results, reporting a significant decrease in shoot Cl content with Si applications. However, Romero-Aranda et al. (2006), and Soylemezoglu et al. (2009), reported a slight increase in Cl shoots concentration with applications of Si.

A significant finding with regards Na content was the reduction of Na within shoots when Si treated plants was exposed to 60 mM NaCl when compared to the control (Table 1). The decrease of Na content in Si treated plants has been a major finding when evaluating the ability of Si to mitigate NaCl salinity stress. Many studies that evaluate Si and NaCl relate NaCl mitigation to a reduction in Na shoot concentrations (Saqib et al., 2008; Tuna et al., 2008; Savvas et al., 2009; Ashraf et al., 2010), the reduction of which may be linked to Si reducing transpiration, as concluded by Ma et al. (2001). Savvas (2007), also reported a decrease in Na with applications of Si, but did not conclude this to be factor influencing

the ability of Si to mitigate salinity stress.

Other mechanisms of salinity mitigation by Si have been suggested, including the ability of Si to improve water storage within plant tissues. Due to the higher growth rate that this would allow, it follows that salt concentrations would decrease through dilution, and therefore ameliorate salt toxicity (Romero-Aranda et al., 2006). Anti oxidants have also been a topic of investigation, with Liang (1999) and Zhu et al. (2004a, b), investigating the interaction of Si with antioxidants in salt stressed barley, cucumber and tomato respectively, with promising results.

Although transpiration and anti oxidants were not factors investigated in this study, plant weight increases, and subsequent decreases in Na shoot concentrations, leads to the conclusion that mitigation of NaCl induced salinity with applications of Si was at least partially due to the reduction of shoot Na content. Additions of Si to plants under salinity stress can be recommended, especially at higher salinity levels, as shown by the results of this paper.

ACKNOWLEDGEMENTS

This study was funded by the Cape Peninsula University of Technology, and supported by the Mauerberger Foundation Scholarship.

REFERENCES

- Ahmad R, Zaheer SH, Ismail S (1992). Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Sci.*, 85(1): 43-50.
- Arnon DL (1949). Copper enzymes in isolated chloroplasts. Polyphenoloxidases in *Beta vulgaris*. *Plant Physiol.*, 24: 1-15.

- Arnon DL, Stout PR (1939). The essentiality of certain elements in minute quantities with special reference to copper. *Plant Physiol.*, 14: 371-375.
- Ashraf M, Rahmatullah, Afzal M, Ahmed R, Mujeeb F, Sarwar A, Ali L (2010). Alleviation of detrimental effects of NaCl by silicon nutrition in salt-sensitive and salt-tolerant genotypes of sugarcane (*Saccharum officinarum* L.). *Plant Soil*, 326(1): 381-391.
- Belanger RR, Bowen PA, Ehret DL, Menzies JG (1995). Soluble silicon: its role in crop and disease management of greenhouse crops. *Plant Dis.*, 79(4): 329-336.
- Bradbury M, Ahmad R (1990). The effect of silicon on the growth of *Prosopis juliflora* growing in saline soil. *Plant Soil*, 125: 71-74.
- Carter DL (1981). Salinity and plant productivity. *CRC Handbook of Agricultural Productivity*, 1: 117-133.
- Epstein E (1994). The anomaly of silicon in plant biology. *Proceedings of the National Academy of Sciences of the United States of America*, 91: 11-17.
- Epstein E (1999). Silicon. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 50: 641-664.
- Epstein E, Bloom AJ (2005). Mineral nutrition of plants: Principles and perspectives. 2nd Ed. Sunderland, MA: Sinauer Associates.
- Hadfield J (2001). The A-Z of vegetable gardening in South Africa. Cape Town: Struik, pp. 110.
- Harris D (1992). Hydroponics: The complete guide to gardening without soil. Cape Town: New Holland Publishing.
- Hiscox JD, Israelstam GF (1979). A method for the extraction of chlorophyll from leaf tissue without maceration. *Can. J. Biol.*, 57: 1332-1334.
- Hashemi A, Abdolzadeh A, Sadeghipour HR (2010). Beneficial effects of silicon nutrition in alleviating salinity stress in hydroponically grown canola, *Brassica napus* L., plants. *Soil Sci. Plant Nutr.*, 56(2): 244-353.
- Liang T (1999). Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. *Plant and Soil*, 209: 217-224.
- Maas EV, Hoffman GJ (1977). Crop salt tolerance – Current Assessment. *J. Irrigat. Drain. Div.*, 103: 115-134.
- Ma JF, Miyake Y, Takahashi E (2001). Silicon as a beneficial element for crop plants. In Datnoff, L.E., Snyder, G.H. and Korndörfer, G.H. (Eds). *Silicon in agriculture*. New York: Elsevier, pp. 17-39.
- Marshak S (2005). *Earth: Portrait of a Planet*. 2nd Ed. New York: W.W. Norton and Company, p. 43.
- Marschner H (1995). *Mineral Nutrition of Higher Plants*. 2nd Ed. London: Academic Press, pp. 417-426.
- Muckle ME (1993). *Hydroponic nutrients: Easy ways to make your own*. Canada: Growers Press.
- Reezi S, Babalar M, Kalantari S (2009). Silicon alleviates salt stress, decreases malondialdehyde content and affects petal color of saltstressed cut rose (*Rosa xhybrida* L.) 'Hot Lady'. *Afr. J. Biotechnol.*, 8(8): 1502-1508.
- Romero-Aranda MR, Jurado O, Cuartero J (2006). Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *J. Plant Physiol.*, 163(8): 847-855.
- Savvas D, Giotis D, Chatzieustratiou D, Bakea M, Patakioutas G (2009). Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environ. Exp. Bot.*, 65(1): 11-17.
- Savvas D, Kizas G, Lydakis-Simantiris N, Salahas G, Papadimitriou M, Tsouka N (2007). Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *Eur. J. Hort. Sci.*, 72(2): 73-79.
- Shannon MC, Grieve CN (1999). Tolerance of vegetables to salinity. *Sci. Hort.*, 78: 5-38.
- Saqib M, Zorb C, Schubert S (2008). Silicon-mediated improvement in the salt resistance of wheat (*Triticum aestivum*) results from increased sodium exclusion and resistance to oxidative stress. *Funct. Plant Biol.*, 35(7): 633-639.
- Soylemezoglu G, Demir K, Inal A, Gunes A (2009). Effect of silicon on antioxidant and stomatal response of two grapevine (*Vitis vinifera* L.) rootstocks grown in boron toxic, saline and boron toxic-saline soil. *Sci. Hort.*, 123(2): 240-246.
- Stern KR (2006). *Introductory plant biology*. 10th Ed. New York: McGraw Hill, p. 152, 161.
- Tas G, Papadandonakis N, Savvas D (2005). Response of lettuce (*Lactuca sativa* L. var. *longifolia*) grown in a closed hydroponic system to NaCl- or CaCl₂-salinity. *J. Appl. Bot. Food. Qual.*, 79(2): 136-140.
- Tuna AL, Kaya C, Higgs D, Murillo-Amador B, Aydemir S, Girgin AR (2008). Silicon improves salinity tolerance in wheat plants. *Environ. Exp. Bot.*, 62(1): 10-16.
- Urdan TC (2005). *Statistics in Plain English*. 2nd Ed. New Jersey: Lawrence Erlbaum Associates, pp. 120-170.
- Wang X, Wei Z, Liu D, Zhao G (2011). Effects of NaCl and silicon on activities of antioxidative enzymes in roots, shoots and leaves of alfalfa. *Afr. J. Biotechnol.*, 10(4): 545-549.
- Zhu Z, Wei G, Li J, Qian Q, Yu J (2004a). Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Sci.*, 167: 527-533.
- Zhu Z, Al-aghabary K, Shi Q (2004b). Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *J. Plant Nutr.*, 27(12): 2101-2115.

Salinity induced changes in oxidative stress and antioxidant status as affected by applications of silicon in lettuce (*Lactuca sativa*)

Christopher J. Milne^a, Charles P. Laubscher^{a,*}, Patrick A. Ndakidemi^b, Jeanine L. Marnewick^c, Fanie Rautenbach^c

^a Department of Horticultural Sciences, Cape Peninsula University of Technology, Cape Town, South Africa

^b The Nelson Mandela African Institute of Science and Technology, Arusha-Tanzania

^c Oxidative Stress Research Centre, Department of Biomedical Sciences, Cape Peninsula University of Technology, Cape Town, South Africa

*Corresponding author. Email: laubscherc@cput.ac.za

Abstract

The antioxidant content, capacity and certain oxidative stress parameters were investigated in salinity stressed *Lactuca sativa* L. 'Eish!' with the addition of exogenous silicon (Si). No significant changes were shown for total polyphenol concentrations in any of the experimental plants. Applications of 4mM Si significantly increased oxygen radical absorbance capacity analysis (ORAC) values at both 30 and 60mM NaCl. The redox status of glutathione was also improved with a significant increase in reduced glutathione (GSH) in 2mM Si treated plants at 30mM NaCl. Increases in both GSH and oxidised glutathione (GSSG) were noted at 60mM NaCl. The activity of catalase (CAT) was

significantly decreased with the addition of 1 and 4mM Si at 30mM NaCl. Lipid peroxidation (measured as thiobarbituric acid reactive substances, TBARS) was not influenced by the different Si concentrations. These results suggest that applications of Si could be beneficial with regards to the modulation of oxidative stress in salinity stressed lettuce, and should be considered as an addition when cultivating lettuce.

Key Words: *Potassium silicate, ROS, Antioxidants, Hydroponics*

Introduction

Salinity is a significant plant stress factor and a major problem on all continents (Ghassemi *et al.*, 1995). Saline conditions influence plant growth in two core ways; by increasing the osmotic potential of the soil solution, which has a direct effect on plant-water availability (Carter, 1981), and through ion toxicities, determined by the specific ions that are in excess (Marschner, 1995). In most cases, sodium (Na^+) and chloride (Cl^-) ions are the cause of saline conditions (Harris, 1992; Epstein & Bloom, 2005). The effect of NaCl induced salinity on plants differ from plant to plant, but generally leads to a reduced growth rate, sometimes nutritional deficiencies, and at higher concentrations, the accumulation of Na^+ and Cl^- , resulting in leaf scorching, chlorosis and necrosis (Shannon & Grieve, 1999; Wahid *et al.*, 1999). Along with these symptoms, salinity stress is known to result in an excess production of reactive oxygen species (ROS) (Jaspers *et al.*, 2005), oxidative damage and a change in

concentrations of antioxidants (Bor *et al.*, 2003; Sekmen *et al.*, 2007; Gao *et al.*, 2008). Consequently, ROS are good cellular indicators of stress (Mittler, 2002).

Silicon (Si), a quasi-essential element for plants (Epstein, 1999), has proved to have significant ameliorative properties regarding NaCl toxicity in many important agricultural and ornamental crops. It has shown to repeatedly result in an increase in yield together with other beneficial growth parameters (Romero-Aranda *et al.*, 2006; Savvas *et al.*, 2007; Soylemezoglu *et al.*, 2009). Silicon allows plants to grow more productively on saline land, leading to definite economic benefits, especially for developing nations (Shay, 1990).

The literature disagrees in the manner by which Si mitigates NaCl toxicity. Si has the ability to influence Na absorption, distribution or transportation (Ahmad *et al.*, 1992; Savvas *et al.*, 2009; Soylemezoglu *et al.*, 2009; Hashemi, *et al.*, 2010), which is sometimes associated with an increase in potassium (K) concentration (Saqib *et al.*, 2008; Tuna *et al.*, 2008; Ashraf *et al.*, 2010). Ameliorative properties of silicon have also been associated with the decrease in plant Na concentrations through dilution, thus mitigating salt toxicity (Romero-Aranda *et al.*, 2006). Ma *et al.* (2001), suggested silicon's beneficial effects stems from the decreased Na influx through a Si induced decrease in transpiration. In addition to Si – Na related amelioration of salinity stress, studies have indicated that Si can influence antioxidant activity. This has been reported in, amongst others, salt stressed barley, (Liang, 1999), cucumber, (Zhu *et al.*, 2004a), tomato (Zhu *et al.*, 2004b) and wheat (Saqib *et al.*, 2008).

In this study the influence of Si in NaCl stressed *Lactuca sativa* L. 'Eish!' was investigated in order to assess whether the known Si induced yield increase can be associated with antioxidant concentrations and oxidative damage. Furthermore, it is to our knowledge that this is the first study examining Si applications influencing antioxidant status and oxidative damage in NaCl salinity stressed lettuce.

Materials and Methods

Plant Conditions and Cultivation

Lettuce was cultivated in a research greenhouse at the Cape Peninsula University of Technology (Cape Town, South Africa). Greenhouse temperatures ranged between a minimum of nine and maximum of 29°C, and humidity between a minimum of 44% and a maximum of 99%. Seeds were sown in a 128 plug polystyrene seedling tray, with vermiculite as the sowing medium. Seeds germinated eight days after sowing, and the seedlings were fertilized daily with half strength NUTRIFEED (Starke Ayres, South Africa); a complete, water soluble fertilizer providing 9.3mM N, 1.7mM P, 6.7mM K, 3.5mM Ca, 1.8mM Mg, 4.7mM S, 53.7µM Fe, 8.7µM Mn, 44.4µM B, 1.5µM Zn, 6.3µM Cu and 0.2µM Mo. Four weeks after sowing, 80 seedlings were chosen at random, and transplanted into 12.5cm plastic pots, with HYDROTON (Germany) expanded clay used as the medium, and then moved directly into the soilless growing systems. There were a total of eight, closed, sub irrigative, soilless growing

systems, delivering ± 1 cm of recirculating nutrient solution. Each of these eight systems had a different treatment, equating to eight treatments with 10 plants (repeats) per treatment.

Basal Nutrient Solution

Salts used to add macro nutrients to the nutrient solution were of analytical grade; KNO_3 , $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, $\text{NH}_4\text{H}_2\text{PO}_4$, and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ were used to include 16mM N, 6 mM K, 4 mM Ca, 2 mM P, 1 mM S and 1 mM Mg respectively, based on the modified Hoagland solution as described by Epstein & Bloom (2005). Micro nutrients were supplied using HYGROPLEX (Hygrotech, South Africa); a complete commercially available micro nutrient mix, adding 58 μM B, 11.4 μM Mn, 7.6 μM Zn, 1 μM Cu, 0.65 μM Mo and 31.9 μM Fe to the nutrient solution. The basal nutrient solution, having had an electrical conductivity (EC) of 1.9 dS m^{-1} , was formulated using deionized water.

Silicon and NaCl Treatments

Si was added to the basal nutrient solution using K_2SiO_3 (AGRISIL K50), at concentrations of 0, 1, 2 and 4mM Si. Two concentrations of NaCl (30 & 60mM), were applied to each of the 4 Si concentrations, making a total of 8 treatments. Treatments were applied in a single dose, implemented one week after transplantation. Newly prepared treatments containing 0, 1, 2 and 4mM Si and 30mM NaCl had a mean EC of 5.20 dS m^{-1} , and treatments containing 0, 1,

2 and 4mM Si and 60mM NaCl, had a mean EC of 8.10 dS m⁻¹. These EC levels were maintained daily. K that was added with the applications of K₂SiO₃ was balanced in all treatments by subtracting additional K from KNO₃, and supplementing lost NO₃ by adding HNO₃. The nutrient solution, which was renewed weekly, was maintained daily at a pH of 6.0 using HCl and NaOH.

Sampling and Harvesting

Antioxidant status and oxidative damage

The shoots of 5 of the repeats per treatment were harvested 6 weeks after being transplanted into the soilless growing system. Once harvested, shoots were rinsed twice in deionized water and patted dry with paper towel. The shoots were then placed directly in an air tight plastic bag, and stored at -80°C. The shoots were transported on dry ice, and lyophilized at -80°C for 16 h. Once the lyophilization was complete, samples were ground to a powder using a mortar and pestle and stored at -80°C until analyzed.

Plant yield and elemental analysis

Plant yield is expressed as fresh and dry weight of roots and shoot in grams. Five plants per treatments were separated into roots and shoots, rinsed twice in deionised water and pattered dry with paper towel. After recording the fresh weights, the plant material were placed into paper bags and dried at 65°C.

Once the weight of the dried plant material stayed constant, dry shoot and root weights were recorded.

Quantities of shoot Si were measured by dissolving the ground, ashed material in HCl. Concentrations, determined using an inductively-coupled plasma emission spectrophotometer, are expressed in mg g^{-1} of dried plant material.

Total protein analysis

For total protein extraction, 125mg of lyophilized plant material was homogenized with 6mL of 25mM HEPES-KOH buffer containing 0.2mM EDTA and 2% PVP (pH 7.8), on ice. The homogenate was centrifuged in 2mL microcentrifuge tubes at 15 000 *g* for 10 minutes at 4°C. The resulting supernatant was transferred to new 2mL microcentrifuge tubes and stored at -80°C until needed.

Protein content was determined using the PIERCE BCA protein assay kit, utilizing the procedures accompanying the kit, with absorbance levels being measured at 562nm. Total protein content (expressed as $\mu\text{g mL}^{-1}$) was used for catalase activity quantification.

Polyphenol Antioxidant Content and Capacity

Total Polyphenol (TP) Analysis

For TP analysis, 125mg of lyophilized plant material was homogenised with 6mL of methanol on ice, and centrifuged in 2mL microcentrifuge tubes at 15 000 *g* for 10 minutes at 4°C. 25µL undiluted supernatant was then transferred, in triplicate, to a 96 well plate containing 125µL of 10 times diluted Folin-Ciocalteu's phenol reagent (0.2N). After 5 minutes, 100 µL of sodium carbonate (7.5%, w/v) was added and the resulting mixture incubated at room temperature for a further two hours. Absorbance values at 765nm were recorded and compared to those of a Gallic acid standard. Results are expressed as mg GAE g⁻¹.

Oxygen Radical Absorbance Capacity Analysis

Extraction followed that of TP, with 12µL of 10 times diluted supernatant being added to 138µL fluorescein and 50µL AAPH, and analyzed, in triplicate, using a fluorescence spectrophotometer until zero fluorescence occurred. The ORAC values were calculated by dividing the sample curve by that of the Trolox standard, and expressed as µmol TE g⁻¹ (Ou *et al.*, 2001).

Catalase Analysis

Catalase (CAT) activity was measured, in triplicate, by placing 20 μ L of the supernatant (obtained from protein analysis, refer 2.5) with 170 μ L of a 50mM Potassium phosphate buffer (pH7.0) and 75 μ L H₂O₂ in a 96 well micro plate. The disappearance of H₂O₂ was measured over a 2 minute period by measuring absorbance levels at 240nm every 15 seconds. Catalase activity was calculated using the extinction coefficient of H₂O₂ and expressed as $\mu\text{mol min}^{-1} \mu\text{g}^{-1}$ protein (Aebi, 1984).

Glutathione Analysis

For GSH extraction, 125mg of lyophilized plant material was homogenised with 6mL of 5% Trichloroacetic Acid buffer, on ice. The homogenate was centrifuged in 2mL microcentrifuge tubes at 15 000 *g* for 10 minutes at 4°C. The resulting supernatant was transferred to new 2mL microcentrifuge tubes and stored at -80°C until needed.

Glutathione concentrations were measured, in triplicate, by placing 50 μ L of 50 times diluted supernatant, with 50 μ L of 0.3mM DTNB (5,5'-Dithiobis(2-nitrobenzoic acid)) and 50 μ L glutathione reductase (GR) solution (0.02U/ μ L) in a 96 well micro plate. After a 5 minute incubation period at room temperature, 50 μ L of 1mM NADPH was placed in each well. Absorbance levels at 412nm were measured at 15 second intervals for 5 minutes. Absorbance levels were

compared to those of a GSH standard curve, and total GSH expressed as $\mu\text{mol g}^{-1}$ dried plant material.

GSSG extraction followed that of GSH, except for the addition of 4.8mg of M2VP (1-Methyl-2-vinylpyridinium) within the homogenate buffer. Measurement of GSSG followed that of GSH, except a 25 times diluted supernatant was used, and absorbance levels were compared to those of a GSSG standard curve (Asensi *et al.*, 1999).

Lipid Peroxidation Analysis

For thiobarbituric acid reactive substances extraction, the method by Yagi, (1984), adapted for plant samples, was utilised. 125mg of lyophilized plant material was homogenised with 6mL of Methanol, on ice. The homogenate was centrifuged in 2mL microcentrifuge tubes at 15 000 *g* for 10 minutes at room temperature. The resulting supernatant was transferred to new 2mL microcentrifuge tubes and stored at -80°C until needed.

250 μL of undiluted supernatant, with 375 μL of 0.44mM PCA (perchloric acid) and 125 μL of 42mM TBA (2-Thiobarbituric acid) were placed in 2mL microcentrifuge tubes. After a 60 minute incubation period at 100°C , the 2mL microcentrifuge tubes were centrifuged at 10 000 *g* for 5 minutes at room temperature. The resulting supernatant was placed, in triplicate, in a 96 well plate, and absorbance at 532nm were compared to those of the

malondialdehyde standard and expressed as $\mu\text{mol g}^{-1}$ dried plant material (Yagi, 1984).

Statistical Analysis

Statistical analysis was performed using two-way analysis of variance (ANOVA), followed by the Bonferroni post test, with P values ≤ 0.05 considered significant. Pearson correlations coefficient was used to calculate correlations. Computations were executed with the software program SPSS (Urdan, 2005).

Results

Plant weights showed no significant differences when treated plants were compared to the 30mM NaCl control. However, there were a 12 and 17% increase in fresh shoot and root weight and a 46% increase in dry shoot weight when comparing the 4mM Si treatment to the control (Table I).

However, plant weights showed a clear increase when treated plants were compared to the 60mM NaCl control. Although 2mM Si was the only treatment that resulted in a marginally significant increase ($P \leq 0.1$) in fresh shoot weight, the increase was considerable (72%). The fresh root weight of the 2mM Si treated plants resulted in a significant increase (75%) when compared to the control. Dry shoot weights significantly increase by 74 and 75% with both 1 and

2mM Si treated plants respectively when compared to the control. Although 4mM Si treated plants showed no significant difference when compared to the 60mM NaCl control, there was a marked increase of 56%. Dry root weights showed a marginally significant increase of 68% with 1mM Si when compared to the 60mM NaCl control. Again, although dry root weights of 2 and 4mM Si treated plants resulted in no significant differences when compared to the control, there was a noteworthy increase of 50 and 43% respectively.

Shoot Si concentrations of 1mM Si treated plants showed no significant difference at both 30 and 60mM NaCl when compared to their individual controls. Nevertheless, treatments of 2 and 4mM Si at 30mM NaCl resulted in a significant increase of 150% and 616% respectively. Similarly, treatments of 2 and 4mM Si at 60mM NaCl resulted in a significant increase of 129% and 157% respectively (Fig. 1).

With regards to antioxidant status and oxidative stress between the two controls, there were no significant differences between any of the parameters, besides that of the TP controls. The 60mM NaCl control showed a 18% increase in TP concentrations when compared to that of the 30mM NaCl control, resulting in a marginal difference ($P \leq 0.1$) between the two controls. However, when Si treated plants were compared to that of the controls, TP concentrations showed no differences (Table II). ORAC values showed a tendency to increase with applications of Si at all concentrations when compared to the controls. However, only 4mM Si at both 30 and 60mM NaCl

showed significant increases when compared to controls, with a 28% and 21% increase respectively.

Catalase activities showed a clear tendency to decrease with applications of Si when plants were exposed to 30mM NaCl, with 1 and 4mM Si, a decrease of 29 and 25% respectively, being significant when compared to the 30mM NaCl control (Table III). Conversely Si applications at 60mM NaCl showed no tendency to decrease, and showed no significance when compared to that of the 60mM control.

Concentrations of both GSH and GSSG, like that of ORAC, tended to increase with applications of Si. Concentrations of GSH at 2mM Si with 30mM NaCl, and both 1 and 4mM Si with 60mM NaCl showed significant increases of 138%, 106% and 99% respectively, when compared to their individual controls. Concentrations of GSSG showed a significant increase of 100% at 4mM Si, and a marginally significance increase of 93% ($P \leq 0.1$) at 1mM Si when compared to the 60mM NaCl control. The ratio of GSH:GSSG showed a limited trend to increase or decrease, with an application of 2mM Si with 30mM NaCl being the only treatment showing a significant increase of 33% ($P \leq 0.1$) when compared to the control.

Like that of TP concentrations, TBARS values showed no significant differences when any of the treatments were compared to their respective controls. In addition to this, no correlations were found when comparing variables within treatments.

Discussion

Plant stresses, including salinity stress, are known to disturb cellular homeostasis, enhancing the production of ROS. (Dat *et al.*, 2000). Additionally, osmotic stress, one of the foremost stresses associated with high salinity levels, has been shown to cause the production of ROS (Xiong & Zhu, 2002).

Although ROS have roles as signalling molecules, active generation of which can be initiated by abiotic stresses (Desikan *et al.*, 2005), ROS are traditionally thought of as species causing cell injury or death (Mittler, 2002). In the light of this, plants need ways to detoxify ROS. CAT is a major ROS scavenging enzyme seen as indispensable, with stress bringing about the production of CAT (Mittler, 2002). With ROS causing the production of CAT, it can be argued that lower concentrations of CAT, indicating lower levels of ROS, would be a sign of less oxidative stress. In this study, with the application of Si, there was an overall tendency for CAT concentrations to decrease, significantly with 1 and 4mM at 30mM NaCl. Zhu *et al.* (2004a), however, showed that Si raises CAT levels in salt treated tomatoes, while Zhu *et al.* (2004b) reported tendencies for CAT levels to decrease with applications of Si on salt treated plants. Bor *et al.* (2003), Gao *et al.* (2008) and Eraslan *et al.* (2007), although not investigating Si, found applications of NaCl to significantly increase CAT activities, supporting the idea that increased CAT activities could be indicative of oxidative stress. Gong *et al.* (2005) when investigating applications of Si to ameliorate drought stress, found that applications of Si lead to significantly decreased CAT activity.

This is relevant to NaCl stress, as osmotic stress is a primary limitation for plants growing in saline conditions (Marschner, 1995).

Contrary to CAT activities, yet similarly indicating a decrease in oxidative stress, ORAC levels showed a tendency to increase, significantly at 4mM Si with both 30 and 60mM NaCl. ORAC, a hydrophilic antioxidant capacity assay, is one of the most commonly used methods for measuring antioxidative capacity in biology (Yeum *et al.*, 2010). It is assumed that with an increase in TP concentrations, there would be a correlative increase in ORAC values. The level of TP in Si treated plants showed no significant differences between controls, and perhaps even a tendency to decrease. Liu *et al.* (2007) reported similar results, although with relatively higher TP concentrations, with no correlation to the DPPH assay, a procedure that has shown to be comparable to that of the ORAC assay (Thaipong *et al.*, 2006), even though different mechanisms are involved (Ndhlala *et al.*, 2010).

TBARS concentrations, a well-known indicator of lipid peroxidation (Wada *et al.*, 2008), a process that can lead to cell death (Mittler, 2002), showed no significant differences between the treated and the control plants, and no trend to increase or decrease. This is in disagreement with Liang (1999), Zhu *et al.* (2004a & b), all finding applications of Si to decrease lipid peroxidation.

Increases in GSH concentrations are known to be associated with salinity stress (Ruiz & Blumwald, 2002; Leyva *et al.*, 2011). The results from this study showed that applications of Si tended to increase GSH concentrations,

significantly with 2mM Si at 30mM NaCl and with 1 and 4mM Si at 60mM NaCl, when compared to NaCl alone. These results are in agreement with Saquib *et al.* (2008), reporting that GSH increases with applications of Si in NaCl stressed wheat. Foyer *et al.* (2005) argue that the signal responses that could be brought about by the interaction between GSH and the GSH:GSSG ratio are linked to plant growth cessation. This is interesting when considering that, contrary to conventional thought, active plant growth cessation, as opposed to stress limiting growth, has been linked to surviving adverse environmental conditions (Harberd *et al.*, 2009), including salinity stress (Magome *et al.*, 2008). However, applications of Si are widely reported to increase plant growth (Romero-Aranda *et al.*, 2006; Soylemezoglu *et al.*, 2009), which is in line with the data presented in this study. Therefore, applications of Si induced increases in GSH concentrations seem unlikely to be associated with a growth cessation stress response. GSH are well known antioxidants, and higher concentrations would infer superior antioxidative defence (Tausz *et al.*, 2004), and would therefore logically result in a decrease in ROS concentrations brought about by salinity stress (Foyer *et al.*, 2005). This approach would arguably provide more probable grounds for the increases in GSH with applications of Si.

The overall impression of Si applications in relation to antioxidant concentrations and oxidative stress modulation ROS is that fertilization and regular inclusion of Si in nutrient solutions, where Si is often limited, is of value to lettuce cultivation. Investigations into the effects of higher salinity concentrations would likely show clear cut results with regards to the effects of Si applications on antioxidant status in salinity stressed lettuce, and is a

recommended focus for future studies. In addition to this, an investigation into other antioxidant related factors such as ascorbate peroxidase, glutathione reductase and ascorbic acid could provide more insight into Si induced amelioration of oxidative stress.

References

- Aebi, H., 1984. Catalase in vitro. *Methods Enzymol.*, **105**: 121–126
- Ahmad, R., S.H. Zaheer and S. Ismail, 1992 .Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Sci.*, **85**: 43-50
- Asensi, M., J. Sastre, F.V. Pallardo, A. Lloret, M. Lehner, J. Garcia-de-la Asuncion and J. Viña, 1999. Ratio of reduced to oxidised glutathione as indicator of oxidative stress status and DNA damage. *Methods Enzymol.*, **299**: 267-76
- Ashraf, M., Rahmatullah, M. Afzal, R. Ahmed, F. Mujeeb, A. Sarwar and L. Ali, 2010. Alleviation of detrimental effects of NaCl by silicon nutrition in salt-sensitive and salt-tolerant genotypes of sugarcane (*Saccharum officinarum* L.). *Plant Soil.*, **326**: 381-391

Bor, M., F. Özdemir and I. Türkan, 2003. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Sci.*, **164**: 77-184

Carter, D.L., 1981. Salinity and plant productivity. CRC Handbook of Agricultural Productivity. **1**: 117-133

Dat, J., S. Vandenabeele, E. Vranová, M. Van Montagu, D. Inzé and F. Van Breusegem, 2000. Dual action of the active oxygen species during plant stress responses. *Cell Mol. Life Sci.*, **57**: 779–795

Desikan, R., K. Hancock and S. Neill, 2005. Reactive oxygen species as signalling molecules. In: Smirnov, N. (ed.). *Antioxidants and Reactive Oxygen Species in Plants*. pp: 169-196. Blackwell Publishing, India

Epstein, E. 1999. Silicon. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **50**: 641-664

Epstein, E. and A.J. Bloom, 2005. *Mineral nutrition of plants: Principles and perspectives*, Sinauer Associates, Massachusetts

Eraslan, F., A. Inal, O. Savasturk and A. Gunes, 2007. Changes in antioxidative system and membrane damage of lettuce in response to salinity and boron toxicity. *Sci. Hort.*, **114**: 5–10

Foyer, C.H., F.L. Gomez and P.D.R. van Heerden, 2005. Glutathione. In: Smirnoff, N. (ed.). *Antioxidants and Reactive Oxygen Species in Plants*. pp. 1 - 24. Blackwell Publishing, India

Gao, S., C. Ouyang, S. Wang, T. Xu, L. Tang and F. Chen, 2008. Effects of salt stress on growth, antioxidant enzyme and phenylalanine ammonia-lyase activities in *Jatropha curcas* L. seedlings. *Plant Soil Environ.*, **54**: 374–381

Ghassemi F., A.J. Jakeman and H.A Hix, 1995. *Salinisation of land and water resources: human causes, extent, management and case studies*. CABI Press, Australia

Gong, H., X. Zhu, K. Chen, S. Wang and C. Zhang, 2005. Silicon alleviates oxidative damage of wheat plants in pots under drought. - *Plant Sci.*, **169**: 313–321

Harberd, N.P., E. Belfield and Y. Yasumura, 2009. The angiosperm gibberellin-GID1-DELLA growth regulatory mechanism: How an "inhibitor of an inhibitor" enables flexible response to fluctuating environments. *Plant Cell.*, **21**: 1328-1339

Harris, D., 1992. *Hydroponics: The complete guide to gardening without soil*. New Holland Publishing, South Africa

- Hashemi, A., A. Abdolzadeh, and H.R. Sadeghipour, 2010. Beneficial effects of silicon nutrition in alleviating salinity stress in hydroponically grown canola, *Brassica napus* L., plants. *Soil Sci. Plant Nutr.*, **56**: 244-253
- Jaspers, P., H. Kollist, C. Langebartels and J. Kangasjärvi, 2005. Plant responses to ozone. - In: Smirnoff, N. (ed.). *Antioxidants and Reactive Oxygen Species in Plants*. pp: 268-292. Blackwell Publishing, India
- Leyva, R., E. Sánchez-Rodríguez, J.J. Ríos, M.M. Rubio-Wilhelmi, J.M. Romero, L., Ruiz and B. Blasco, 2011. Beneficial effects of exogenous iodine in lettuce plants subjected to salinity stress. *Plant Sci.*, **181**: 195–202
- Liang, T., 1999. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barley under salt stress. *Plant Soil.*, **209**: 217-224
- Liu, X., S. Ardo, M. Bunning, J. Parry, K. Zhou, C. Stushnoff, F. Stoniker and L. Yu, 2007. Total phenolics content and DPPH radical scavenging activity of lettuce (*Lactuca sativa* L.) grown in Colorado. *LWT - Food Sci. Technol.*, **40**: 552-557
- Ma, J.F., Y. Miyake and E. Takahashi, 2001. Silicon as a beneficial element for crop plants. In: Datnoff, L.E., G.H. Snyder and G.H. Korndörfer (eds.). *Silicon in agriculture*. pp. 17-39. Elsevier, New York

- Magome, H., S. Yamaguchi, A. Hanada, Y. Kamiya and K. Oda, 2008. The DDF1 transcriptional activator upregulates expression of a gibberellin-deactivating gene, *GA2ox7*, under high-salinity stress in *Arabidopsis*. *The Plant J.* **56**: 613-626
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. Academic Press, London
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.*, **7**: 405-410
- Ndhkala, A.R., M. Moyo and J. Van Staden 2010. Natural antioxidants: Fascinating or mythical biomolecules? *Molecules.*, **15**: 6905-6930
- Ou, B., M. Hampsch-Woodill and R.L. Prior, 2001. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J. Agric. Food Chem.*, **49**: 4619-4626
- Romero-Aranda, M.R., O. Jurado and J. Cuartero, 2006. Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *J. Plant Physiol.*, **163**: 847-855
- Ruiz, J.M. and E. Blumwald, 2002. Salinity-induced glutathione synthesis in *Brassica napus*. *Planta.*, **214**: 965–969

- Saqib, M., C. Zorb and S. Schubert, 2008. Silicon-mediated improvement in the salt resistance of wheat (*Triticum aestivum*) results from increased sodium exclusion and resistance to oxidative stress. *Funct. Plant Biol.*, **35**: 633-639
- Savvas, D., D. Giotis, D. Chatzieustratiou, M. Bakea and G. Patakioutas, 2009. Silicon supply in soilless cultivations of zucchini alleviates stress induced by salinity and powdery mildew infections. *Environ. Exp. Bot.*, **65**: 11-17
- Savvas, D., G. Kizas, N. Lydakis-Simantiris, G. Salahas, M. Papadimitriou and N. Tsouka, 2007. Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *European. J. Hort. Sci.*, **72**: 73-79
- Sekmen, A.H., I. Türkana and S. Takiob, 2007. Differential responses of antioxidative enzymes and lipid peroxidation to salt stress in salt-tolerant *Plantago maritime* and salt-sensitive *Plantago media*. *Physiol. Plant.*, **131**: 399–411
- Shannon, M.C., and C.N. Grieve, 1999. Tolerance of vegetables to salinity. *Sci. Hort.*, **78**, 5-38
- Shay, G., 1990. *Saline Agriculture: Salt Tolerant Plants for Developing Countries*. National Academy Press, Washington
- Soylemezoglu, G., K. Demir, A. Inal and A. Gunes, 2009. Effect of silicon on antioxidant and stomatal response of two grapevine (*Vitis vinifera* L.)

rootstocks grown in boron toxic, saline and boron toxic-saline soil. *Sci. Hort.*, **123**: 240-246

Tausz, M., H. Sircelj and D. Grill, 2004. The glutathione system as a stress marker in plant ecophysiology: is a stress response concept valid? *J. Exp. Bot.*, **55**: 1855-1862

Thaipong, K., U. Boonprakob, K. Crosby, L. Cisneros-Zevallos and D. Byrne, 2006. Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *J. Food Comp. Anal.*, **19**: 669–675

Tuna, A.L., C. Kaya, D. Higgs, B. Murillo-Amador, S. Aydemir and A.R. Girgin, 2008. Silicon improves salinity tolerance in wheat plants. *Environ. Exp. Bot.*, **62**: 10-16

Urdan, T.C., 2005. *Statistics in Plain English*. Lawrence Erlbaum Associates, New Jersey

Wada. M., N. Kishikawa, N. Kuroda and K. Nakashima, 2008. Evaluation method for antioxidative activity of health food. In: Papadopoulos, K.N. (ed.). *Food Chemistry Research Developments*. pp: 199-221. Nova Science Publishers, New York

- Wahid, A., I. Masood, I-ul-H. Javed and E. Rasul. 1999. Phenotypic flexibility as marker of sodium chloride tolerance in sunflower genotypes. *Environ. Exp. Bot.*, 42: 85–94
- Xiong, L. and J.K. Zhu, 2002. Molecular and genetic aspects of plant response to osmotic stress. *Plant Cell Environ.*, **25**: 131–139
- Yagi, K., 1984. Assay for plasma lipid peroxides. *Methods Enzymol.*, **109**: 328-331
- Yeum, K., R. Russell and G. Aldini, 2010. Antioxidant activity and oxidative stress: An overview. In: Aldini, G., K. Yeum, E. Niki and R. Russell (eds.). *Biomarkers for antioxidant defense and oxidative damage*. pp: 3-20. Blackwell Publishing, Iowa
- Zhu, Z., G. Wei, J. Li, Q. Qian and J. Yu, 2004a. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Sci.*, **167**: 527–533
- Zhu, Z., K. Al-aghabary and Q. Shi, 2004b. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *J. Plant Nutr.*, **27**: 2101-211

Table I:

Effects of Si applications on plant fresh and dry root and shoot weights in salinity stressed lettuce.

NaCl	Si	Fresh Weight (g)		Dry Weight (g)	
		Shoots	Roots	Shoots	Roots
30mM	0mM	47.85 ± 17.56	6.49 ± 1.83	2.04 ± 0.51	0.42 ± 0.1
	1mM	39.1 ± 13.07	5.74 ± 1.81	1.94 ± 0.73	0.34 ± 0.1
	2mM	43.52 ± 10.49	5.75 ± 1.54	2.12 ± 0.55	0.36 ± 0.09
	4mM	53.70 ± 11.54	7.64 ± 1.60	2.99 ± 0.56	0.44 ± 0.09
60mM	0mM	28.05 ± 6.55	4.52 ± 1.27	1.47 ± 0.31	0.28 ± 0.1
	1mM	41.73 ± 12.39	7.29 ± 2.38	2.56 ± 0.73**	0.47 ± 0.19*
	2mM	48.12 ± 14.92*	7.92 ± 2.06**	2.57 ± 0.69**	0.42 ± 0.14
	4mM	40.97 ± 9.83	6.44 ± 1.52	2.29 ± 0.59	0.40 ± 0.05

Mean values of treated plants were compared to that of the controls (0mM Si). Significance levels are represented by *P < 0.1. **P < 0.05.

Table II:
Effects of Si applications on Total Polyphenolics (TP) and Oxidative Radical Absorbance Capacity, in salinity stressed lettuce.

NaCl	Si	TP (mg GAE g ⁻¹)	ORAC (μmol TE g ⁻¹)
30mM	0mM	5.05 ± 0.37	49.28 ± 3.93
	1mM	4.23 ± 0.35	56.81 ± 4.23
	2mM	4.71 ± 0.59	54.99 ± 4.07
	4mM	5.19 ± 0.63	63.16 ± 5.83*
60mM	0mM	5.97 ± 0.58	47.72 ± 3.22
	1mM	5.25 ± 0.59	55.12 ± 9.86
	2mM	4.98 ± 0.50	54.70 ± 5.74
	4mM	5.85 ± 0.65	57.51 ± 2.79**

Mean values of treated plants were compared to that of the controls (0mM Si). Significance levels are represented by *P ≤ 0.1, **P ≤ 0.05, ***P ≤ 0.001.

Table III:

Effects of Si applications on Thiobarbituric Acid Reactive Substances (TBARS), Glutathione (GSH), Glutathione disulfide (GSSG), GSH:GSSG ratio and Catalase (CAT), in salinity stressed lettuce.

NaCl	Si	TBARS ($\mu\text{M g}^{-1}$)	GSH ($\mu\text{mol g}^{-1}$)	GSSG ($\mu\text{mol g}^{-1}$)	GSH:GSSG	CAT ($\mu\text{mol min}^{-1} \text{ug}^{-1}$)
30mM	0mM	0.26 \pm 0.03	0.91 \pm 0.37	0.15 \pm 0.06	5.96 \pm 0.81	1.03 \pm 0.09
	1mM	0.26 \pm 0.04	0.99 \pm 0.2	0.19 \pm 0.05	5.20 \pm 0.27	0.73 \pm 0.09***
	2mM	0.24 \pm 0.03	2.17 \pm 0.24***	0.23 \pm 0.08	7.94 \pm 0.59*	0.86 \pm 0.12
	4mM	0.29 \pm 0.04	1.43 \pm 0.46	0.24 \pm 0.07	5.91 \pm 0.67	0.77 \pm 0.11**
60mM	0mM	0.25 \pm 0.02	1.04 \pm 0.83	0.15 \pm 0.1	6.94 \pm 1.76	1.12 \pm 0.10
	1mM	0.28 \pm 0.03	2.14 \pm 0.37**	0.29 \pm 0.07*	7.69 \pm 1.87	1.03 \pm 0.06
	2mM	0.25 \pm 0.03	1.47 \pm 0.38	0.25 \pm 0.06	5.87 \pm 0.15	0.95 \pm 0.11
	4mM	0.26 \pm 0.03	2.07 \pm 0.28**	0.30 \pm 0.05**	7.06 \pm 1.01	1.28 \pm 0.16

Mean values of treated plants were compared to that of the controls (0mM Si). Significance levels are represented by *P \leq 0.1, **P \leq 0.05, ***P \leq 0.001.




Figure 1. Effect of Si applications on Si shoot concentrations in salinity stressed lettuce.